M^{censed}^{TO}LTIVARIABLE CALCULUS

LARSON EDWARDS

NINTH EDITION



Multivariable Calculus, Ninth Edition Larson/Edwards

VP/Editor-in-Chief: Michelle Julet Publisher: Richard Stratton Senior Sponsoring Editor: Cathy Cantin Development Editor: Peter Galuardi Associate Editor: Jeannine Lawless Editorial Assistant: Amy Haines Media Editor: Peter Galuardi Senior Content Manager: Maren Kunert Senior Marketing Manager: Jennifer Jones Marketing Communications Manager: Mary Anne Payumo Senior Content Project Manager, Editorial Production: Tamela Ambush Art and Design Manager: Jill Haber Senior Manufacturing Coordinator: Diane Gibbons Permissions Editor: Katie Huha Text Designer: Nesbitt Graphics Art Editor: Larson Texts, Inc. Senior Photo Editor: Jennifer Meyer Dare Illustrator: Larson Texts, Inc. Cover Designer: Harold Burch Cover Image: © Richard Edelman/Woodstock Graphics Studio Compositor: Larson Texts, Inc.

TI is a registered trademark of Texas Instruments, Inc. Mathematica is a registered trademark of Wolfram Research, Inc. Maple is a registered trademark of Waterloo Maple, Inc.

Problems from the William Lowell Putnam Mathematical Competition reprinted with permission from the Mathematical Association of America, 1529 Eighteenth Street, NW. Washington, DC. © 2010, 2006 Brooks/Cole, Cengage Learning

ALL RIGHTS RESERVED. No part of this work covered by the copyright herein may be reproduced, transmitted, stored or used in any form or by any means graphic, electronic, or mechanical, including but not limited to photocopying, recording, scanning, digitizing, taping, Web distribution, information networks, or information storage and retrieval systems, except as permitted under Section 107 or 108 of the 1976 United States Copyright Act, without the prior written permission of the publisher.

For product information and technology assistance, contact us at Cengage Learning Customer & Sales Support, 1-800-354-9706

For permission to use material from this text or product, submit all requests online at **www.cengage.com/permissions**. Further permissions questions can be emailed to **permissionrequest@cengage.com**

Library of Congress Control Number: 2008939232 Student Edition: ISBN-13: 978-0-547-20997-5 ISBN-10: 0-547-20997-5

Brooks/Cole

10 Davis Drive Belmont, CA 94002-3098 USA

Cengage learning is a leading provider of customized learning solutions with office locations around the globe, including Singapore, the United Kingdom, Australia, Mexico, Brazil, and Japan. Locate your local office at: **international.cengage.com/region**

Cengage learning products are represented in Canada by Nelson Education, Ltd.

For your course and learning solutions, visit www.cengage.com

Purchase any of our products at your local college store or at our preferred online store **www.ichapters.com**

Printed in the United States of America 1 2 3 4 5 6 7 12 11 10 09 08

Vectors and the Geometry of Space

This chapter introduces vectors and the three-dimensional coordinate system. Vectors are used to represent lines and planes, and are also used to represent quantities such as force and velocity. The three-dimensional coordinate system is used to represent surfaces such as ellipsoids and elliptical cones. Much of the material in the remaining chapters relies on an understanding of this system.

In this chapter, you should learn the following.

- How to write vectors, perform basic vector operations, and represent vectors graphically. (11.1)
- How to plot points in a three-dimensional coordinate system and analyze vectors in space. (11.2)
- How to find the dot product of two vectors (in the plane or in space). (11.3)
- How to find the cross product of two vectors (in space). (11.4)
- How to find equations of lines and planes in space, and how to sketch their graphs. (11.5)
- How to recognize and write equations of cylindrical and quadric surfaces and of surfaces of revolution. (11.6)
- How to use cylindrical and spherical coordinates to represent surfaces in space. (11.7)



Mark Hunt/Hunt Stock

Two tugboats are pushing an ocean liner, as shown above. Each boat is exerting a force of 400 pounds. What is the resultant force on the ocean liner? (See Section 11.1, Example 7.)



Vectors indicate quantities that involve both magnitude and direction. In Chapter 11, you will study operations of vectors in the plane and in space. You will also learn how to represent vector operations geometrically. For example, the graphs shown above represent vector addition in the plane.

764

11.1 Vectors in the Plane



A directed line segment Figure 11.1



Equivalent directed line segments **Figure 11.2**





The vectors **u** and **v** are equivalent. **Figure 11.3**

- Write the component form of a vector.
- Perform vector operations and interpret the results geometrically.
- Write a vector as a linear combination of standard unit vectors.
- Use vectors to solve problems involving force or velocity.

Component Form of a Vector

Many quantities in geometry and physics, such as area, volume, temperature, mass, and time, can be characterized by a single real number scaled to appropriate units of measure. These are called **scalar quantities**, and the real number associated with each is called a **scalar**.

Other quantities, such as force, velocity, and acceleration, involve both magnitude and direction and cannot be characterized completely by a single real number. A **directed line segment** is used to represent such a quantity, as shown in Figure 11.1. The directed line segment \overrightarrow{PQ} has **initial point** P and **terminal point** Q, and its **length** (or **magnitude**) is denoted by $\|\overrightarrow{PQ}\|$. Directed line segments that have the same length and direction are **equivalent**, as shown in Figure 11.2. The set of all directed line segments that are equivalent to a given directed line segment \overrightarrow{PQ} is a **vector in the plane** and is denoted by $\mathbf{v} = \overrightarrow{PQ}$. In typeset material, vectors are usually denoted by lowercase, boldface letters such as \mathbf{u} , \mathbf{v} , and \mathbf{w} . When written by hand, however, vectors are often denoted by letters with arrows above them, such as \overrightarrow{u} , \overrightarrow{v} , and \overrightarrow{w} .

Be sure you understand that a vector represents a *set* of directed line segments (each having the same length and direction). In practice, however, it is common not to distinguish between a vector and one of its representatives.

EXAMPLE 1 Vector Representation by Directed Line Segments

Let **v** be represented by the directed line segment from (0, 0) to (3, 2), and let **u** be represented by the directed line segment from (1, 2) to (4, 4). Show that **v** and **u** are equivalent.

Solution Let P(0, 0) and Q(3, 2) be the initial and terminal points of **v**, and let R(1, 2) and S(4, 4) be the initial and terminal points of **u**, as shown in Figure 11.3. You can use the Distance Formula to show that \overline{PQ} and \overline{RS} have the *same length*.

$$\begin{aligned} |\overline{PQ}|| &= \sqrt{(3-0)^2 + (2-0)^2} = \sqrt{13} & \text{Length of } \overline{PQ} \\ ||\overline{RS}|| &= \sqrt{(4-1)^2 + (4-2)^2} = \sqrt{13} & \text{Length of } \overline{RS} \end{aligned}$$

Both line segments have the *same direction*, because they both are directed toward the upper right on lines having the same slope.

Slope of
$$\overrightarrow{PQ} = \frac{2-0}{3-0} = \frac{2}{3}$$

and

Slope of
$$\overrightarrow{RS} = \frac{4-2}{4-1} = \frac{2}{3}$$

Because \overrightarrow{PQ} and \overrightarrow{RS} have the same length and direction, you can conclude that the two vectors are equivalent. That is, **v** and **u** are equivalent.



A vector in standard position **Figure 11.4**

The directed line segment whose initial point is the origin is often the most convenient representative of a set of equivalent directed line segments such as those shown in Figure 11.3. This representation of **v** is said to be in **standard position**. A directed line segment whose initial point is the origin can be uniquely represented by the coordinates of its terminal point $Q(v_1, v_2)$, as shown in Figure 11.4.

DEFINITION OF COMPONENT FORM OF A VECTOR IN THE PLANE

If **v** is a vector in the plane whose initial point is the origin and whose terminal point is (v_1, v_2) , then the **component form of v** is given by

 $\mathbf{v} = \langle v_1, v_2 \rangle.$

The coordinates v_1 and v_2 are called the **components of v.** If both the initial point and the terminal point lie at the origin, then **v** is called the **zero vector** and is denoted by $\mathbf{0} = \langle 0, 0 \rangle$.

This definition implies that two vectors $\mathbf{u} = \langle u_1, u_2 \rangle$ and $\mathbf{v} = \langle v_1, v_2 \rangle$ are **equal** if and only if $u_1 = v_1$ and $u_2 = v_2$.

The following procedures can be used to convert directed line segments to component form or vice versa.

If P(p₁, p₂) and Q(q₁, q₂) are the initial and terminal points of a directed line segment, the component form of the vector v represented by PQ is ⟨v₁, v₂⟩ = ⟨q₁ - p₁, q₂ - p₂⟩. Moreover, from the Distance Formula you can see that the length (or magnitude) of v is

$$\|\mathbf{v}\| = \sqrt{(q_1 - p_1)^2 + (q_2 - p_2)^2}$$
Length of a vector

$$= \sqrt{v_1^2 + v_2^2}.$$

2. If $\mathbf{v} = \langle v_1, v_2 \rangle$, \mathbf{v} can be represented by the directed line segment, in standard position, from P(0, 0) to $Q(v_1, v_2)$.

The length of **v** is also called the **norm of v**. If $||\mathbf{v}|| = 1$, **v** is a **unit vector**. Moreover, $||\mathbf{v}|| = 0$ if and only if **v** is the zero vector **0**.

EXAMPLE 2 Finding the Component Form and Length of a Vector

Find the component form and length of the vector **v** that has initial point (3, -7) and terminal point (-2, 5).

Solution Let $P(3, -7) = (p_1, p_2)$ and $Q(-2, 5) = (q_1, q_2)$. Then the components of $\mathbf{v} = \langle v_1, v_2 \rangle$ are

$$w_1 = q_1 - p_1 = -2 - 3 = -5$$

 $w_2 = q_2 - p_2 = 5 - (-7) = 12$

So, as shown in Figure 11.5, $\mathbf{v} = \langle -5, 12 \rangle$, and the length of \mathbf{v} is

$$\|\mathbf{v}\| = \sqrt{(-5)^2 + 12^2} \\ = \sqrt{169} \\ = 13.$$

Q(-2,5) 6 4 -6 -4 -2 -2 -4 -6 -4 -6 -8P(3,-7)

Component form of v: $\mathbf{v} = \langle -5, 12 \rangle$ Figure 11.5



The scalar multiplication of v Figure 11.6



WILLIAM ROWAN HAMILTON (1805 - 1865)

Some of the earliest work with vectors was done by the Irish mathematician William Rowan Hamilton. Hamilton spent many years developing a system of vector-like quantities called quaternions. Although Hamilton was convinced of the benefits of quaternions, the operations he defined did not produce good models for physical phenomena. It wasn't until the latter half of the nineteenth century that the Scottish physicist James Maxwell (1831–1879) restructured Hamilton's quaternions in a form useful for representing physical quantities such as force, velocity, and acceleration.

Vector Operations

DEFINITIONS OF VECTOR ADDITION AND SCALAR MULTIPLICATION

- Let $\mathbf{u} = \langle u_1, u_2 \rangle$ and $\mathbf{v} = \langle v_1, v_2 \rangle$ be vectors and let *c* be a scalar.
- **1.** The vector sum of **u** and **v** is the vector $\mathbf{u} + \mathbf{v} = \langle u_1 + v_1, u_2 + v_2 \rangle$.
- **2.** The scalar multiple of c and u is the vector $c\mathbf{u} = \langle cu_1, cu_2 \rangle$.
- 3. The **negative** of **v** is the vector

$$-\mathbf{v} = (-1)\mathbf{v} = \langle -v_1, -v_2 \rangle.$$

4. The difference of **u** and **v** is

$$\mathbf{u} - \mathbf{v} = \mathbf{u} + (-\mathbf{v}) = \langle u_1 - v_1, u_2 - v_2 \rangle.$$

Geometrically, the scalar multiple of a vector **v** and a scalar c is the vector that is |c| times as long as v, as shown in Figure 11.6. If c is positive, cv has the same direction as **v**. If c is negative, c**v** has the opposite direction.

The sum of two vectors can be represented geometrically by positioning the vectors (without changing their magnitudes or directions) so that the initial point of one coincides with the terminal point of the other, as shown in Figure 11.7. The vector $\mathbf{u} + \mathbf{v}$, called the **resultant vector**, is the diagonal of a parallelogram having \mathbf{u} and v as its adjacent sides.



Figure 11.8 shows the equivalence of the geometric and algebraic definitions of vector addition and scalar multiplication, and presents (at far right) a geometric interpretation of $\mathbf{u} - \mathbf{v}$.



Figure 11.8



EXAMPLE 3 Vector Operations

Given $\mathbf{v} = \langle -2, 5 \rangle$ and $\mathbf{w} = \langle 3, 4 \rangle$, find each of the vectors. **a.** $\frac{1}{2}\mathbf{v}$ **b.** $\mathbf{w} - \mathbf{v}$ **c.** $\mathbf{v} + 2\mathbf{w}$ Solution **a.** $\frac{1}{2}\mathbf{v} = \langle \frac{1}{2}(-2), \frac{1}{2}(5) \rangle = \langle -1, \frac{5}{2} \rangle$ **b.** $\mathbf{w} - \mathbf{v} = \langle w_1 - v_1, w_2 - v_2 \rangle = \langle 3 - (-2), 4 - 5 \rangle = \langle 5, -1 \rangle$ **c.** Using $2\mathbf{w} = \langle 6, 8 \rangle$, you have $\mathbf{v} + 2\mathbf{w} = \langle -2, 5 \rangle + \langle 6, 8 \rangle$ $= \langle -2 + 6, 5 + 8 \rangle$ $= \langle 4, 13 \rangle$.

Vector addition and scalar multiplication share many properties of ordinary arithmetic, as shown in the following theorem.

THEOREM 11.1 PROPERTIES OF VECTOR OPERATIONS			
Let \mathbf{u} , \mathbf{v} , and \mathbf{w} be vectors in the plane, and let c and d be scalars.			
1. $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$ Commutative Property			
2. $(u + v) + w = u + (v + w)$	Associative Property		
3. $u + 0 = u$	Additive Identity Property		
4. $\mathbf{u} + (-\mathbf{u}) = 0$	Additive Inverse Property		
5. $c(d\mathbf{u}) = (cd)\mathbf{u}$			
$6. \ (c+d)\mathbf{u} = c\mathbf{u} + d\mathbf{u}$	Distributive Property		
7. $c(\mathbf{u} + \mathbf{v}) = c\mathbf{u} + c\mathbf{v}$	Distributive Property		
8. $1(\mathbf{u}) = \mathbf{u}, 0(\mathbf{u}) = 0$			

PROOF The proof of the *Associative Property* of vector addition uses the Associative Property of addition of real numbers.

$$(\mathbf{u} + \mathbf{v}) + \mathbf{w} = [\langle u_1, u_2 \rangle + \langle v_1, v_2 \rangle] + \langle w_1, w_2 \rangle$$

= $\langle u_1 + v_1, u_2 + v_2 \rangle + \langle w_1, w_2 \rangle$
= $\langle (u_1 + v_1) + w_1, (u_2 + v_2) + w_2 \rangle$
= $\langle u_1 + (v_1 + w_1), u_2 + (v_2 + w_2) \rangle$
= $\langle u_1, u_2 \rangle + \langle v_1 + w_1, v_2 + w_2 \rangle = \mathbf{u} + (\mathbf{v} + \mathbf{w})$

Similarly, the proof of the *Distributive Property* of vectors depends on the Distributive Property of real numbers.

$$(c + d)\mathbf{u} = (c + d)\langle u_1, u_2 \rangle$$

= $\langle (c + d)u_1, (c + d)u_2 \rangle$
= $\langle cu_1 + du_1, cu_2 + du_2 \rangle$
= $\langle cu_1, cu_2 \rangle + \langle du_1, du_2 \rangle = c\mathbf{u} + d\mathbf{u}$

The other properties can be proved in a similar manner.

Copyright 2010 Cengage Learning, Inc. All Rights Reserved. May not be copied, scanned, or duplicated, in whole or in part.

Bchap



EMMY NOETHER (1882-1935)

One person who contributed to our knowledge of axiomatic systems was the German mathematician Emmy Noether. Noether is generally recognized as the leading woman mathematician in recent history.

FOR FURTHER INFORMATION For

more information on Emmy Noether, see the article "Emmy Noether, Greatest Woman Mathematician" by Clark Kimberling in *The Mathematics Teacher*. To view this article, go to the website *www.matharticles.com*.

lichar

Any set of vectors (with an accompanying set of scalars) that satisfies the eight properties given in Theorem 11.1 is a **vector space**.* The eight properties are the *vector space axioms*. So, this theorem states that the set of vectors in the plane (with the set of real numbers) forms a vector space.

THEOREM 11.2 LENGTH OF A SCALAR MULTIPLE

Let **v** be a vector and let c be a scalar. Then

 $\|c\mathbf{v}\| = |c| \|\mathbf{v}\|.$ |c| is the absolute value of c.

PROOF Because
$$c\mathbf{v} = \langle cv_1, cv_2 \rangle$$
, it follows that

$$\begin{aligned} \|c\mathbf{v}\| &= \|\langle cv_1, cv_2 \rangle\| = \sqrt{(cv_1)^2 + (cv_2)^2} \\ &= \sqrt{c^2v_1^2 + c^2v_2^2} \\ &= \sqrt{c^2(v_1^2 + v_2^2)} \\ &= |c|\sqrt{v_1^2 + v_2^2} \\ &= |c| \|\mathbf{v}\|. \end{aligned}$$

In many applications of vectors, it is useful to find a unit vector that has the same direction as a given vector. The following theorem gives a procedure for doing this.

THEOREM 11.3 UNIT VECTOR IN THE DIRECTION OF v

If \mathbf{v} is a nonzero vector in the plane, then the vector

$$\mathbf{u} = \frac{\mathbf{v}}{\|\mathbf{v}\|} = \frac{1}{\|\mathbf{v}\|}\mathbf{v}$$

has length 1 and the same direction as v.

PROOF Because $1/||\mathbf{v}||$ is positive and $\mathbf{u} = (1/||\mathbf{v}||)\mathbf{v}$, you can conclude that \mathbf{u} has the same direction as \mathbf{v} . To see that $||\mathbf{u}|| = 1$, note that

$$\|\mathbf{u}\| = \left\| \left(\frac{1}{\|\mathbf{v}\|}\right) \mathbf{v} \right\|$$
$$= \left| \frac{1}{\|\mathbf{v}\|} \right\| \|\mathbf{v}\|$$
$$= \frac{1}{\|\mathbf{v}\|} \|\mathbf{v}\|$$
$$= 1$$

So, **u** has length 1 and the same direction as **v**.

In Theorem 11.3, **u** is called a **unit vector in the direction of v.** The process of multiplying **v** by $1/||\mathbf{v}||$ to get a unit vector is called **normalization of v.**

^{*} For more information about vector spaces, see Elementary Linear Algebra, Sixth Edition, by Larson and Falvo (Boston: Houghton Mifflin Harcourt Publishing Company, 2009).

EXAMPLE 4 Finding a Unit Vector

Find a unit vector in the direction of $\mathbf{v} = \langle -2, 5 \rangle$ and verify that it has length 1.

Solution From Theorem 11.3, the unit vector in the direction of **v** is

$$\frac{\mathbf{v}}{\|\mathbf{v}\|} = \frac{\langle -2, 5 \rangle}{\sqrt{(-2)^2 + (5)^2}} = \frac{1}{\sqrt{29}} \langle -2, 5 \rangle = \left\langle \frac{-2}{\sqrt{29}}, \frac{5}{\sqrt{29}} \right\rangle.$$

This vector has length 1, because

$$\sqrt{\left(\frac{-2}{\sqrt{29}}\right)^2 + \left(\frac{5}{\sqrt{29}}\right)^2} = \sqrt{\frac{4}{29} + \frac{25}{29}} = \sqrt{\frac{29}{29}} = 1.$$

Generally, the length of the sum of two vectors is not equal to the sum of their lengths. To see this, consider the vectors \mathbf{u} and \mathbf{v} as shown in Figure 11.9. By considering \mathbf{u} and \mathbf{v} as two sides of a triangle, you can see that the length of the third side is $\|\mathbf{u} + \mathbf{v}\|$, and you have

$$\|\mathbf{u} + \mathbf{v}\| \le \|\mathbf{u}\| + \|\mathbf{v}\|$$

Equality occurs only if the vectors \mathbf{u} and \mathbf{v} have the *same direction*. This result is called the **triangle inequality** for vectors. (You are asked to prove this in Exercise 91, Section 11.3.)

Standard Unit Vectors

The unit vectors $\langle 1, 0 \rangle$ and $\langle 0, 1 \rangle$ are called the **standard unit vectors** in the plane and are denoted by

$$\mathbf{i} = \langle 1, 0 \rangle$$
 and $\mathbf{j} = \langle 0, 1 \rangle$ Standard unit vectors

as shown in Figure 11.10. These vectors can be used to represent any vector uniquely, as follows.

$$\mathbf{v} = \langle v_1, v_2 \rangle = \langle v_1, 0 \rangle + \langle 0, v_2 \rangle = v_1 \langle 1, 0 \rangle + v_2 \langle 0, 1 \rangle = v_1 \mathbf{i} + v_2 \mathbf{j}$$

The vector $\mathbf{v} = v_1 \mathbf{i} + v_2 \mathbf{j}$ is called a **linear combination** of \mathbf{i} and \mathbf{j} . The scalars v_1 and v_2 are called the **horizontal** and **vertical components of v.**

EXAMPLE 5 Writing a Linear Combination of Unit Vectors

Let **u** be the vector with initial point (2, -5) and terminal point (-1, 3), and let $\mathbf{v} = 2\mathbf{i} - \mathbf{j}$. Write each vector as a linear combination of **i** and **j**.

a. u b. w = 2u - 3v

Solution

a.
$$\mathbf{u} = \langle q_1 - p_1, q_2 - p_2 \rangle$$

= $\langle -1 - 2, 3 - (-5) \rangle$
= $\langle -3, 8 \rangle = -3\mathbf{i} + 8\mathbf{j}$
b. $\mathbf{w} = 2\mathbf{u} - 3\mathbf{v} = 2(-3\mathbf{i} + 8\mathbf{j}) - 3(2\mathbf{i} - \mathbf{j})$
= $-6\mathbf{i} + 16\mathbf{j} - 6\mathbf{i} + 3\mathbf{j}$
= $-12\mathbf{i} + 19\mathbf{j}$



Triangle inequality Figure 11.9



Standard unit vectors i and j Figure 11.10

770 Chapter 11 Vectors and the Geometry of Space



The angle θ from the positive *x*-axis to the vector **u** Figure 11.11

If **u** is a unit vector and θ is the angle (measured counterclockwise) from the positive *x*-axis to **u**, then the terminal point of **u** lies on the unit circle, and you have

$$\mathbf{u} = \langle \cos \theta, \sin \theta \rangle = \cos \theta \mathbf{i} + \sin \theta \mathbf{j}$$
 Unit vector

as shown in Figure 11.11. Moreover, it follows that any other nonzero vector **v** making an angle θ with the positive *x*-axis has the same direction as **u**, and you can write

 $\mathbf{v} = \|\mathbf{v}\| \langle \cos \theta, \sin \theta \rangle = \|\mathbf{v}\| \cos \theta \mathbf{i} + \|\mathbf{v}\| \sin \theta \mathbf{j}.$

EXAMPLE 6 Writing a Vector of Given Magnitude and Direction

The vector **v** has a magnitude of 3 and makes an angle of $30^\circ = \pi/6$ with the positive *x*-axis. Write **v** as a linear combination of the unit vectors **i** and **j**.

Solution Because the angle between v and the positive x-axis is $\theta = \pi/6$, you can write the following.

$$= \|\mathbf{v}\| \cos \theta \mathbf{i} + \|\mathbf{v}\| \sin \theta \mathbf{j}$$

= $3 \cos \frac{\pi}{6} \mathbf{i} + 3 \sin \frac{\pi}{6} \mathbf{j}$
= $\frac{3\sqrt{3}}{2} \mathbf{i} + \frac{3}{2} \mathbf{j}$

Applications of Vectors

v

Vectors have many applications in physics and engineering. One example is force. A vector can be used to represent force, because force has both magnitude and direction. If two or more forces are acting on an object, then the **resultant force** on the object is the vector sum of the vector forces.

EXAMPLE 7 Finding the Resultant Force

Two tugboats are pushing an ocean liner, as shown in Figure 11.12. Each boat is exerting a force of 400 pounds. What is the resultant force on the ocean liner?

Solution Using Figure 11.12, you can represent the forces exerted by the first and second tugboats as

- $\mathbf{F}_1 = 400 \langle \cos 20^\circ, \sin 20^\circ \rangle$
 - $= 400 \cos(20^\circ)\mathbf{i} + 400 \sin(20^\circ)\mathbf{j}$
- $\mathbf{F}_2 = 400 \langle \cos(-20^\circ), \sin(-20^\circ) \rangle$
 - $= 400 \cos(20^\circ)\mathbf{i} 400 \sin(20^\circ)\mathbf{j}.$

The resultant force on the ocean liner is

 $\mathbf{F} = \mathbf{F}_1 + \mathbf{F}_2$

 $= [400 \cos(20^\circ)\mathbf{i} + 400 \sin(20^\circ)\mathbf{j}] + [400 \cos(20^\circ)\mathbf{i} - 400 \sin(20^\circ)\mathbf{j}]$ $= 800 \cos(20^\circ)\mathbf{i} \approx 752\mathbf{i}.$

So, the resultant force on the ocean liner is approximately 752 pounds in the direction of the positive *x*-axis.

In surveying and navigation, a **bearing** is a direction that measures the acute angle that a path or line of sight makes with a fixed north-south line. In air navigation, bearings are measured in degrees clockwise from north.



The resultant force on the ocean liner that is exerted by the two tugboats Figure 11.12







(b) Direction with wind **Figure 11.13**

11.1 Exercises

EXAMPLE 8 Finding a Velocity

An airplane is traveling at a fixed altitude with a negligible wind factor. The airplane is traveling at a speed of 500 miles per hour with a bearing of 330° , as shown in Figure 11.13(a). As the airplane reaches a certain point, it encounters wind with a velocity of 70 miles per hour in the direction N 45° E (45° east of north), as shown in Figure 11.13(b). What are the resultant speed and direction of the airplane?

Solution Using Figure 11.13(a), represent the velocity of the airplane (alone) as

 $\mathbf{v}_1 = 500 \cos(120^\circ)\mathbf{i} + 500 \sin(120^\circ)\mathbf{j}.$

The velocity of the wind is represented by the vector

 $\mathbf{v}_2 = 70 \cos(45^\circ)\mathbf{i} + 70 \sin(45^\circ)\mathbf{j}.$

See www.CalcChat.com for worked-out solutions to odd-numbered exercises

The resultant velocity of the airplane (in the wind) is

$$= \mathbf{v}_1 + \mathbf{v}_2 = 500 \cos(120^\circ)\mathbf{i} + 500 \sin(120^\circ)\mathbf{j} + 70 \cos(45^\circ)\mathbf{i} + 70 \sin(45^\circ)\mathbf{j}$$

\$\approx - 200.5\mathbf{i} + 482.5\mathbf{j}.

To find the resultant speed and direction, write $\mathbf{v} = \|\mathbf{v}\|(\cos\theta\,\mathbf{i} + \sin\theta\,\mathbf{j})$. Because $\|\mathbf{v}\| \approx \sqrt{(-200.5)^2 + (482.5)^2} \approx 522.5$, you can write

$$\mathbf{v} \approx 522.5 \left(\frac{-200.5}{522.5} \mathbf{i} + \frac{482.5}{522.5} \mathbf{j} \right) \approx 522.5 [\cos(112.6^\circ)\mathbf{i} + \sin(112.6^\circ)\mathbf{j}].$$

The new speed of the airplane, as altered by the wind, is approximately 522.5 miles per hour in a path that makes an angle of 112.6° with the positive *x*-axis.

In Exercises 1–4, (a) find the component form of the vector v and (b) sketch the vector with its initial point at the origin.



In Exercises 5-8, find the vectors u and v whose initial and terminal points are given. Show that u and v are equivalent.

5. u: (3, 2), (5, 6)	6. u: $(-4, 0)$, $(1, 8)$
v : (1, 4), (3, 8)	v : (2, -1), (7, 7)
7. u : (0, 3), (6, −2)	8. u : (−4, −1), (11, −4)
v : (3, 10), (9, 5)	v : (10, 13), (25, 10)

In Exercises 9–16, the initial and terminal points of a vector v are given. (a) Sketch the given directed line segment, (b) write the vector in component form, (c) write the vector as the linear combination of the standard unit vectors i and j, and (d) sketch the vector with its initial point at the origin.

Initial Point	Terminal Point	Initial Point	Terminal Point
. (2, 0)	(5, 5)	10. (4, -6)	(3, 6)
. (8, 3)	(6, -1)	12. (0, -4)	(-5, -1)

The icon \bigcirc indicates that you will find a CAS Investigation on the book's website. The CAS Investigation is a collaborative exploration of this example using the computer algebra systems Maple and Mathematica.

v

Copyright 2010 Cengage Learning, Inc. All Rights Reserved. May not be copied, scanned, or duplicated, in whole or in part.

9 11

Initial Point	Terminal Point	Initial Point	Terminal Point
13. (6, 2)	(6, 6)	14. (7, −1)	(-3, -1)
15. $\left(\frac{3}{2}, \frac{4}{3}\right)$	$\left(\frac{1}{2},3\right)$	16. (0.12, 0.60)	(0.84, 1.25)

In Exercises 17 and 18, sketch each scalar multiple of v.

17. $\mathbf{v} = \langle 3, 5 \rangle$ (a) $2\mathbf{v}$ (b) $-3\mathbf{v}$ (c) $\frac{7}{2}\mathbf{v}$ (d) $\frac{2}{3}\mathbf{v}$ **18.** $\mathbf{v} = \langle -2, 3 \rangle$ (a) $4\mathbf{v}$ (b) $-\frac{1}{2}\mathbf{v}$ (c) $0\mathbf{v}$ (d) $-6\mathbf{v}$

In Exercises 19–22, use the figure to sketch a graph of the vector. To print an enlarged copy of the graph, go to the website *www.mathgraphs.com*.



19. ·	-u	20.	2 u
21. u	$\mathbf{u} - \mathbf{v}$	22.	$\mathbf{u} + 2\mathbf{v}$

In Exercises 23 and 24, find (a) $\frac{2}{3}u$, (b) v - u, and (c) 2u + 5v.

23. $\mathbf{u} = \langle 4, 9 \rangle$	24. $\mathbf{u} = \langle -3, -8 \rangle$
$\mathbf{v} = \langle 2, -5 \rangle$	$\mathbf{v} = \langle 8, 25 \rangle$

In Exercises 25–28, find the vector v where $u = \langle 2, -1 \rangle$ and $w = \langle 1, 2 \rangle$. Illustrate the vector operations geometrically.

25.	$\mathbf{v} = \frac{3}{2}\mathbf{u}$	26. $v = u + w$
27.	$\mathbf{v} = \mathbf{u} + 2\mathbf{w}$	28. $v = 5u - 3w$

In Exercises 29 and 30, the vector v and its initial point are given. Find the terminal point.

29. $\mathbf{v} = \langle -1, 3 \rangle$; Initial point: (4, 2) **30.** $\mathbf{v} = \langle 4, -9 \rangle$; Initial point: (5, 3)

In Exercises 31–36, find the magnitude of v.

31. $v = 7i$	32. $v = -3i$
33. $\mathbf{v} = \langle 4, 3 \rangle$	34. $\mathbf{v} = \langle 12, -5 \rangle$
35. $v = 6i - 5j$	36. $v = -10i + 3j$

In Exercises 37–40, find the unit vector in the direction of v and verify that it has length 1.

37. $\mathbf{v} = \langle 3, 12 \rangle$	38. $\mathbf{v} = \langle -5, 15 \rangle$
39. $\mathbf{v} = \left< \frac{3}{2}, \frac{5}{2} \right>$	40. $\mathbf{v} = \langle -6.2, 3.4 \rangle$

In Exercises 41-44, find the following.

(a)	u	$\mathbf{(b)} \ \mathbf{v} \ $	(c) $\ \mathbf{u} + \mathbf{v}\ $
(d)	$\left\ \begin{array}{c} \mathbf{u} \\ \ \mathbf{u} \ \end{array} \right\ $	(e) $\left\ \frac{\mathbf{v}}{\ \mathbf{v}\ } \right\ $	(f) $\left\ \frac{\mathbf{u} + \mathbf{v}}{\ \mathbf{u} + \mathbf{v}\ } \right\ $
41.	$\mathbf{u} = \langle 1, -1 \rangle$		42. $\mathbf{u} = \langle 0, 1 \rangle$
	$\mathbf{v} = \langle -1, 2 \rangle$		$\mathbf{v} = \langle 3, -3 \rangle$
43.	$\mathbf{u} = \left\langle 1, \frac{1}{2} \right\rangle$		44. $\mathbf{u} = \langle 2, -4 \rangle$
	$\mathbf{v} = \langle 2, 3 \rangle$		$\mathbf{v} = \langle 5, 5 \rangle$

In Exercises 45 and 46, sketch a graph of u, v, and u + v. Then demonstrate the triangle inequality using the vectors u and v.

45.
$$\mathbf{u} = \langle 2, 1 \rangle$$
, $\mathbf{v} = \langle 5, 4 \rangle$ **46.** $\mathbf{u} = \langle -3, 2 \rangle$, $\mathbf{v} = \langle 1, -2 \rangle$

In Exercises 47–50, find the vector v with the given magnitude and the same direction as u.

Magnitude	Direction
47. $\ \mathbf{v}\ = 6$	$\mathbf{u} = \langle 0, 3 \rangle$
48. $\ \mathbf{v}\ = 4$	$\mathbf{u} = \langle 1, 1 \rangle$
49. $\ \mathbf{v}\ = 5$	$\mathbf{u} = \langle -1, 2 \rangle$
50. $\ \mathbf{v}\ = 2$	$\mathbf{u} = \langle \sqrt{3}, 3 \rangle$

In Exercises 51-54, find the component form of v given its magnitude and the angle it makes with the positive x-axis.

51.	$\ \mathbf{v}\ =3,$	$\theta = 0^{\circ}$	52. $\ \mathbf{v}\ = 5$,	$\theta = 120^{\circ}$
53.	$\ \mathbf{v}\ = 2,$	$\theta = 150^{\circ}$	54. $\ \mathbf{v}\ = 4$,	$\theta = 3.5^{\circ}$

In Exercises 55–58, find the component form of u + v given the lengths of u and v and the angles that u and v make with the positive *x*-axis.

55. $\ \mathbf{u}\ = 1$,	$\theta_{\mathbf{u}} = 0^{\circ}$	56. $ \mathbf{u} = 4$,	$\theta_{\mathbf{u}} = 0^{\circ}$
$\ \mathbf{v}\ =3,$	$\theta_{\rm v} = 45^{\circ}$	$\ \mathbf{v}\ =2,$	$\theta_{\rm v} = 60^{\circ}$
57. $\ \mathbf{u}\ = 2$,	$\theta_{\mathbf{u}} = 4$	58. $ \mathbf{u} = 5$,	$\theta_{\mathbf{u}} = -0.5$
$\ \mathbf{v}\ = 1,$	$\theta_{\rm v} = 2$	$\ \mathbf{v}\ =5,$	$\theta_{\rm v} = 0.5$

WRITING ABOUT CONCEPTS

- **59.** In your own words, state the difference between a scalar and a vector. Give examples of each.
- **60.** Give geometric descriptions of the operations of addition of vectors and multiplication of a vector by a scalar.
- **61.** Identify the quantity as a scalar or as a vector. Explain your reasoning.
 - (a) The muzzle velocity of a gun
 - (b) The price of a company's stock
- **62.** Identify the quantity as a scalar or as a vector. Explain your reasoning.
 - (a) The air temperature in a room
 - (b) The weight of a car

In Exercises 63–68, find a and b such that y = au + bw, where $rac{1}{12}$ In Exercises 79 and 80, use a graphing utility to find the $\mathbf{u} = \langle \mathbf{1}, \mathbf{2} \rangle$ and $\mathbf{w} = \langle \mathbf{1}, -\mathbf{1} \rangle$.

63. $\mathbf{v} = \langle 2, 1 \rangle$	64. $\mathbf{v} = \langle 0, 3 \rangle$
65. $\mathbf{v} = \langle 3, 0 \rangle$	66. $\mathbf{v} = \langle 3, 3 \rangle$
67. $\mathbf{v} = \langle 1, 1 \rangle$	68. $\mathbf{v} = \langle -1, 7 \rangle$

In Exercises 69-74, find a unit vector (a) parallel to and (b) perpendicular to the graph of f at the given point. Then sketch the graph of f and sketch the vectors at the given point.

Function	Point
69. $f(x) = x^2$	(3, 9)
70. $f(x) = -x^2 + 5$	(1, 4)
71. $f(x) = x^3$	(1, 1)
72. $f(x) = x^3$	(-2, -8)
73. $f(x) = \sqrt{25 - x^2}$	(3, 4)
74. $f(x) = \tan x$	$\left(\frac{\pi}{4}, 1\right)$

In Exercises 75 and 76, find the component form of v given the magnitudes of u and u + v and the angles that u and u + vmake with the positive x-axis.

75. $\ \mathbf{u}\ = 1, \ \theta = 45^{\circ}$	76. $\ \mathbf{u}\ = 4, \ \theta = 30^{\circ}$
$\ \mathbf{u} + \mathbf{v}\ = \sqrt{2}, \theta = 90^{\circ}$	$\ \mathbf{u} + \mathbf{v}\ = 6, \theta = 120^{\circ}$

77. *Programming* You are given the magnitudes of **u** and **v** and the angles that **u** and **v** make with the positive *x*-axis. Write a program for a graphing utility in which the output is the following.

- (a) **u**+**v** (b) $\| \mathbf{u} + \mathbf{v} \|$
- (c) The angle that $\mathbf{u} + \mathbf{v}$ makes with the positive x-axis
- (d) Use the program to find the magnitude and direction of the resultant of the vectors shown.



CAPSTONE

- **78.** The initial and terminal points of vector \mathbf{v} are (3, -4) and (9, 1), respectively.
 - (a) Write v in component form.
 - (b) Write \mathbf{v} as the linear combination of the standard unit vectors i and j.
 - (c) Sketch v with its initial point at the origin.
 - (d) Find the magnitude of v.

magnitude and direction of the resultant of the vectors.



81. Resultant Force Forces with magnitudes of 500 pounds and 200 pounds act on a machine part at angles of 30° and -45° , respectively, with the x-axis (see figure). Find the direction and magnitude of the resultant force.



Figure for 81

Figure for 82

- **82.** Numerical and Graphical Analysis Forces with magnitudes of 180 newtons and 275 newtons act on a hook (see figure). The angle between the two forces is θ degrees.
 - (a) If $\theta = 30^{\circ}$, find the direction and magnitude of the resultant force.
 - (b) Write the magnitude M and direction α of the resultant force as functions of θ , where $0^{\circ} \leq \theta \leq 180^{\circ}$.
 - (c) Use a graphing utility to complete the table.

θ	0°	30°	60°	90°	120°	150°	180°
M							
α							

- (d) Use a graphing utility to graph the two functions M and α .
- (e) Explain why one of the functions decreases for increasing values of θ whereas the other does not.
- 83. Resultant Force Three forces with magnitudes of 75 pounds, 100 pounds, and 125 pounds act on an object at angles of 30° , 45° , and 120° , respectively, with the positive x-axis. Find the direction and magnitude of the resultant force.
- 84. Resultant Force Three forces with magnitudes of 400 newtons, 280 newtons, and 350 newtons act on an object at angles of -30° , 45° , and 135° , respectively, with the positive *x*-axis. Find the direction and magnitude of the resultant force.
- 85. Think About It Consider two forces of equal magnitude acting on a point.
 - (a) If the magnitude of the resultant is the sum of the magnitudes of the two forces, make a conjecture about the angle between the forces.

- (b) If the resultant of the forces is **0**, make a conjecture about the angle between the forces.
- (c) Can the magnitude of the resultant be greater than the sum of the magnitudes of the two forces? Explain.
- 86. *Graphical Reasoning* Consider two forces $\mathbf{F}_1 = \langle 20, 0 \rangle$ and $\mathbf{F}_2 = 10 \langle \cos \theta, \sin \theta \rangle$.
 - (a) Find $\|\mathbf{F}_1 + \mathbf{F}_2\|$.
- (b) Determine the magnitude of the resultant as a function of θ . Use a graphing utility to graph the function for $0 \le \theta < 2\pi$.
 - (c) Use the graph in part (b) to determine the range of the function. What is its maximum and for what value of θ does it occur? What is its minimum and for what value of θ does it occur?
 - (d) Explain why the magnitude of the resultant is never 0.
- **87.** Three vertices of a parallelogram are (1, 2), (3, 1), and (8, 4). Find the three possible fourth vertices (see figure).



88. Use vectors to find the points of trisection of the line segment with endpoints (1, 2) and (7, 5).

Cable Tension In Exercises 89 and 90, use the figure to determine the tension in each cable supporting the given load.



- **91.** *Projectile Motion* A gun with a muzzle velocity of 1200 feet per second is fired at an angle of 6° above the horizontal. Find the vertical and horizontal components of the velocity.
- **92.** *Shared Load* To carry a 100-pound cylindrical weight, two workers lift on the ends of short ropes tied to an eyelet on the top center of the cylinder. One rope makes a 20° angle away from the vertical and the other makes a 30° angle (see figure).
 - (a) Find each rope's tension if the resultant force is vertical.
 - (b) Find the vertical component of each worker's force.





Figure for 93

- **93.** *Navigation* A plane is flying with a bearing of 302°. Its speed with respect to the air is 900 kilometers per hour. The wind at the plane's altitude is from the southwest at 100 kilometers per hour (see figure). What is the true direction of the plane, and what is its speed with respect to the ground?
- **94.** *Navigation* A plane flies at a constant groundspeed of 400 miles per hour due east and encounters a 50-mile-per-hour wind from the northwest. Find the airspeed and compass direction that will allow the plane to maintain its groundspeed and eastward direction.

True or False? In Exercises 95–100, determine whether the statement is true or false. If it is false, explain why or give an example that shows it is false.

- **95.** If **u** and **v** have the same magnitude and direction, then **u** and **v** are equivalent.
- **96.** If **u** is a unit vector in the direction of **v**, then $\mathbf{v} = \|\mathbf{v}\| \mathbf{u}$.
- **97.** If $\mathbf{u} = a\mathbf{i} + b\mathbf{j}$ is a unit vector, then $a^2 + b^2 = 1$.
- **98.** If $\mathbf{v} = a\mathbf{i} + b\mathbf{j} = \mathbf{0}$, then a = -b.
- **99.** If a = b, then $||a\mathbf{i} + b\mathbf{j}|| = \sqrt{2}a$.
- 100. If **u** and **v** have the same magnitude but opposite directions, then $\mathbf{u} + \mathbf{v} = \mathbf{0}$.
- **101.** Prove that $\mathbf{u} = (\cos \theta)\mathbf{i} (\sin \theta)\mathbf{j}$ and $\mathbf{v} = (\sin \theta)\mathbf{i} + (\cos \theta)\mathbf{j}$ are unit vectors for any angle θ .
- **102.** *Geometry* Using vectors, prove that the line segment joining the midpoints of two sides of a triangle is parallel to, and one-half the length of, the third side.
- **103.** *Geometry* Using vectors, prove that the diagonals of a parallelogram bisect each other.
- **104.** Prove that the vector $\mathbf{w} = \|\mathbf{u}\|\mathbf{v} + \|\mathbf{v}\|\mathbf{u}$ bisects the angle between \mathbf{u} and \mathbf{v} .
- **105.** Consider the vector $\mathbf{u} = \langle x, y \rangle$. Describe the set of all points (x, y) such that $\|\mathbf{u}\| = 5$.

PUTNAM EXAM CHALLENGE

106. A coast artillery gun can fire at any angle of elevation between 0° and 90° in a fixed vertical plane. If air resistance is neglected and the muzzle velocity is constant $(= v_0)$, determine the set *H* of points in the plane and above the horizontal which can be hit.

This problem was composed by the Committee on the Putnam Prize Competition. \circledcirc The Mathematical Association of America. All rights reserved.

11.2 Space Coordinates and Vectors in Space



The three-dimensional coordinate system Figure 11.14

- Understand the three-dimensional rectangular coordinate system.
- Analyze vectors in space.
- Use three-dimensional vectors to solve real-life problems.

Coordinates in Space

Up to this point in the text, you have been primarily concerned with the two-dimensional coordinate system. Much of the remaining part of your study of calculus will involve the three-dimensional coordinate system.

Before extending the concept of a vector to three dimensions, you must be able to identify points in the **three-dimensional coordinate system.** You can construct this system by passing a z-axis perpendicular to both the x- and y-axes at the origin. Figure 11.14 shows the positive portion of each coordinate axis. Taken as pairs, the axes determine three coordinate planes: the xy-plane, the xz-plane, and the *yz*-plane. These three coordinate planes separate three-space into eight octants. The first octant is the one for which all three coordinates are positive. In this threedimensional system, a point P in space is determined by an ordered triple (x, y, z)where *x*, *y*, and *z* are as follows.

- x = directed distance from yz-plane to P
- y = directed distance from *xz*-plane to *P*
- z = directed distance from xy-plane to P

Several points are shown in Figure 11.15.



Points in the three-dimensional coordinate system are represented by ordered triples. Figure 11.15



Right-handed system Figure 11.16



A three-dimensional coordinate system can have either a left-handed or a righthanded orientation. To determine the orientation of a system, imagine that you are standing at the origin, with your arms pointing in the direction of the positive x- and y-axes, and with the z-axis pointing up, as shown in Figure 11.16. The system is right-handed or left-handed depending on which hand points along the x-axis. In this text, you will work exclusively with the right-handed system.

NOTE The three-dimensional rotatable graphs that are available in the premium eBook for this text will help you visualize points or objects in a three-dimensional coordinate system.



The distance between two points in space **Figure 11.17**



Figure 11.18



$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2}$$
 Distance Formula

EXAMPLE 1 Finding the Distance Between Two Points in Space

The distance between the points (2, -1, 3) and (1, 0, -2) is



A **sphere** with center at (x_0, y_0, z_0) and radius *r* is defined to be the set of all points (x, y, z) such that the distance between (x, y, z) and (x_0, y_0, z_0) is *r*. You can use the Distance Formula to find the **standard equation of a sphere** of radius *r*, centered at (x_0, y_0, z_0) . If (x, y, z) is an arbitrary point on the sphere, the equation of the sphere is

$$(x - x_0)^2 + (y - y_0)^2 + (z - z_0)^2 = r^2$$
 Equation of sphere

as shown in Figure 11.18. Moreover, the midpoint of the line segment joining the points (x_1, y_1, z_1) and (x_2, y_2, z_2) has coordinates

$$\frac{x_2}{2}, \frac{y_1 + y_2}{2}, \frac{z_1 + z_2}{2}$$
). Midpo

Midpoint Formula

EXAMPLE 2 Finding the Equation of a Sphere

Find the standard equation of the sphere that has the points (5, -2, 3) and (0, 4, -3) as endpoints of a diameter.

Solution Using the Midpoint Formula, the center of the sphere is

$$\left(\frac{5+0}{2}, \frac{-2+4}{2}, \frac{3-3}{2}\right) = \left(\frac{5}{2}, 1, 0\right).$$
 Midpoint Formula

By the Distance Formula, the radius is

$$r = \sqrt{\left(0 - \frac{5}{2}\right)^2 + (4 - 1)^2 + (-3 - 0)^2} = \sqrt{\frac{97}{4}} = \frac{\sqrt{97}}{2}$$

Therefore, the standard equation of the sphere is

$$\left(x - \frac{5}{2}\right)^2 + (y - 1)^2 + z^2 = \frac{97}{4}.$$
 Equation of sphere



The standard unit vectors in space **Figure 11.19**



Vectors in Space

In space, vectors are denoted by ordered triples $\mathbf{v} = \langle v_1, v_2, v_3 \rangle$. The **zero vector** is denoted by $\mathbf{0} = \langle 0, 0, 0 \rangle$. Using the unit vectors $\mathbf{i} = \langle 1, 0, 0 \rangle$, $\mathbf{j} = \langle 0, 1, 0 \rangle$, and $\mathbf{k} = \langle 0, 0, 1 \rangle$ in the direction of the positive *z*-axis, the **standard unit vector notation** for **v** is

$$\mathbf{v} = v_1 \mathbf{i} + v_2 \mathbf{j} + v_3 \mathbf{k}$$

as shown in Figure 11.19. If **v** is represented by the directed line segment from $P(p_1, p_2, p_3)$ to $Q(q_1, q_2, q_3)$, as shown in Figure 11.20, the component form of **v** is given by subtracting the coordinates of the initial point from the coordinates of the terminal point, as follows.

$$\mathbf{v} = \langle v_1, v_2, v_3 \rangle = \langle q_1 - p_1, q_2 - p_2, q_3 - p_3 \rangle$$

VECTORS IN SPACE

Let $\mathbf{u} = \langle u_1, u_2, u_3 \rangle$ and $\mathbf{v} = \langle v_1, v_2, v_3 \rangle$ be vectors in space and let *c* be a scalar.

- **1.** Equality of Vectors: $\mathbf{u} = \mathbf{v}$ if and only if $u_1 = v_1, u_2 = v_2$, and $u_3 = v_3$.
- **2.** Component Form: If v is represented by the directed line segment from $P(p_1, p_2, p_3)$ to $Q(q_1, q_2, q_3)$, then

$$\mathbf{v} = \langle v_1, v_2, v_3 \rangle = \langle q_1 - p_1, q_2 - p_2, q_3 - p_3 \rangle.$$

- **3.** Length: $\|\mathbf{v}\| = \sqrt{v_1^2 + v_2^2 + v_3^2}$
- **4.** Unit Vector in the Direction of **v**: $\frac{\mathbf{v}}{\|\mathbf{v}\|} = \left(\frac{1}{\|\mathbf{v}\|}\right) \langle v_1, v_2, v_3 \rangle, \quad \mathbf{v} \neq \mathbf{0}$
- **5.** Vector Addition: $\mathbf{v} + \mathbf{u} = \langle v_1 + u_1, v_2 + u_2, v_3 + u_3 \rangle$
- **6.** Scalar Multiplication: $c\mathbf{v} = \langle cv_1, cv_2, cv_3 \rangle$

NOTE The properties of vector addition and scalar multiplication given in Theorem 11.1 are also valid for vectors in space.

EXAMPLE 3 Finding the Component Form of a Vector in Space

Find the component form and magnitude of the vector **v** having initial point (-2, 3, 1) and terminal point (0, -4, 4). Then find a unit vector in the direction of **v**.

Solution The component form of **v** is

$$\mathbf{v} = \langle q_1 - p_1, q_2 - p_2, q_3 - p_3 \rangle = \langle 0 - (-2), -4 - 3, 4 - 1 \rangle$$
$$= \langle 2, -7, 3 \rangle$$

which implies that its magnitude is

$$\|\mathbf{v}\| = \sqrt{2^2 + (-7)^2 + 3^2} = \sqrt{62}$$

The unit vector in the direction of \mathbf{v} is

$$\mathbf{u} = \frac{\mathbf{v}}{\|\mathbf{v}\|} = \frac{1}{\sqrt{62}} \langle 2, -7, 3 \rangle = \left\langle \frac{2}{\sqrt{62}}, \frac{-7}{\sqrt{62}}, \frac{3}{\sqrt{62}} \right\rangle.$$





Parallel vectors Figure 11.21

Recall from the definition of scalar multiplication that positive scalar multiples of a nonzero vector **v** have the same direction as **v**, whereas negative multiples have the direction opposite of **v**. In general, two nonzero vectors **u** and **v** are **parallel** if there is some scalar *c* such that $\mathbf{u} = c\mathbf{v}$.

DEFINITION OF PARALLEL VECTORS

Two nonzero vectors **u** and **v** are **parallel** if there is some scalar *c* such that $\mathbf{u} = c\mathbf{v}$.

For example, in Figure 11.21, the vectors \mathbf{u} , \mathbf{v} , and \mathbf{w} are parallel because $\mathbf{u} = 2\mathbf{v}$ and $\mathbf{w} = -\mathbf{v}$.

EXAMPLE 4 Parallel Vectors

Vector **w** has initial point (2, -1, 3) and terminal point (-4, 7, 5). Which of the following vectors is parallel to **w**?

a.
$$\mathbf{u} = \langle 3, -4, -1 \rangle$$

b. $\mathbf{v} = \langle 12, -16, 4 \rangle$

Solution Begin by writing **w** in component form.

 $\mathbf{w} = \langle -4 - 2, 7 - (-1), 5 - 3 \rangle = \langle -6, 8, 2 \rangle$

- **a.** Because $\mathbf{u} = \langle 3, -4, -1 \rangle = -\frac{1}{2} \langle -6, 8, 2 \rangle = -\frac{1}{2} \mathbf{w}$, you can conclude that \mathbf{u} is parallel to \mathbf{w} .
- **b.** In this case, you want to find a scalar *c* such that

$$\langle 12, -16, 4 \rangle = c \langle -6, 8, 2 \rangle.$$

$$12 = -6c \rightarrow c = -2$$

$$-16 = 8c \rightarrow c = -2$$

$$4 = -2c \rightarrow c = -2$$

Because there is no c for which the equation has a solution, the vectors are not parallel.

EXAMPLE 5 Using Vectors to Determine Collinear Points

Determine whether the points P(1, -2, 3), Q(2, 1, 0), and R(4, 7, -6) are collinear.

Solution The component forms of \overrightarrow{PQ} and \overrightarrow{PR} are

$$\overrightarrow{PQ} = \langle 2 - 1, 1 - (-2), 0 - 3 \rangle = \langle 1, 3, -3 \rangle$$

and

$$\overrightarrow{PR} = \langle 4 - 1, 7 - (-2), -6 - 3 \rangle = \langle 3, 9, -9 \rangle.$$

These two vectors have a common initial point. So, *P*, *Q*, and *R* lie on the same line if and only if \overrightarrow{PQ} and \overrightarrow{PR} are parallel—which they are because $\overrightarrow{PR} = 3 \overrightarrow{PQ}$, as shown in Figure 11.22.



The points *P*, *Q*, and *R* lie on the same line. Figure 11.22

EXAMPLE 6 Standard Unit Vector Notation

- **a.** Write the vector $\mathbf{v} = 4\mathbf{i} 5\mathbf{k}$ in component form.
- **b.** Find the terminal point of the vector $\mathbf{v} = 7\mathbf{i} \mathbf{j} + 3\mathbf{k}$, given that the initial point is P(-2, 3, 5).

Solution

a. Because **j** is missing, its component is 0 and

 $\mathbf{v} = 4\mathbf{i} - 5\mathbf{k} = \langle 4, 0, -5 \rangle.$

b. You need to find $Q(q_1, q_2, q_3)$ such that $\mathbf{v} = \overrightarrow{PQ} = 7\mathbf{i} - \mathbf{j} + 3\mathbf{k}$. This implies that $q_1 - (-2) = 7$, $q_2 - 3 = -1$, and $q_3 - 5 = 3$. The solution of these three equations is $q_1 = 5$, $q_2 = 2$, and $q_3 = 8$. Therefore, Q is (5, 2, 8).

Application

EXAMPLE 7 Measuring Force

A television camera weighing 120 pounds is supported by a tripod, as shown in Figure 11.23. Represent the force exerted on each leg of the tripod as a vector.

Solution Let the vectors \mathbf{F}_1 , \mathbf{F}_2 , and \mathbf{F}_3 represent the forces exerted on the three legs. From Figure 11.23, you can determine the directions of \mathbf{F}_1 , \mathbf{F}_2 , and \mathbf{F}_3 to be as follows.

$$\overline{PQ}_{1} = \langle 0 - 0, -1 - 0, 0 - 4 \rangle = \langle 0, -1, -4 \rangle$$

$$\overline{PQ}_{2} = \left\langle \frac{\sqrt{3}}{2} - 0, \frac{1}{2} - 0, 0 - 4 \right\rangle = \left\langle \frac{\sqrt{3}}{2}, \frac{1}{2}, -4 \right\rangle$$

$$\overline{PQ}_{3} = \left\langle -\frac{\sqrt{3}}{2} - 0, \frac{1}{2} - 0, 0 - 4 \right\rangle = \left\langle -\frac{\sqrt{3}}{2}, \frac{1}{2}, -4 \right\rangle$$

Because each leg has the same length, and the total force is distributed equally among the three legs, you know that $\|\mathbf{F}_1\| = \|\mathbf{F}_2\| = \|\mathbf{F}_3\|$. So, there exists a constant c such that

$$\mathbf{F}_1 = c\langle 0, -1, -4 \rangle, \quad \mathbf{F}_2 = c \left\langle \frac{\sqrt{3}}{2}, \frac{1}{2}, -4 \right\rangle, \text{ and } \mathbf{F}_3 = c \left\langle -\frac{\sqrt{3}}{2}, \frac{1}{2}, -4 \right\rangle.$$

Let the total force exerted by the object be given by $\mathbf{F} = \langle 0, 0, -120 \rangle$. Then, using the fact that

$$\mathbf{F} = \mathbf{F}_1 + \mathbf{F}_2 + \mathbf{F}_3$$

F

you can conclude that $\mathbf{F}_1, \mathbf{F}_2$, and \mathbf{F}_3 all have a vertical component of -40. This implies that c(-4) = -40 and c = 10. Therefore, the forces exerted on the legs can be represented by

$$\mathbf{F}_1 = \langle 0, -10, -40 \rangle$$

$$\mathbf{F}_2 = \langle 5\sqrt{3}, 5, -40 \rangle$$

$$\mathbf{F}_3 = \langle -5\sqrt{3}, 5, -40 \rangle.$$



780 Chapter 11 Vectors and the Geometry of Space

1.2 Exercises

In Exercises 1 and 2, approximate the coordinates of the points.



In Exercises 3–6, plot the points on the same three-dimensional coordinate system.

3. (a) (2, 1, 3)	(b) $(-1, 2, 1)$
4. (a) (3, −2, 5)	(b) $\left(\frac{3}{2}, 4, -2\right)$
5. (a) (5, −2, 2)	(b) $(5, -2, -2)$
6. (a) (0, 4, −5)	(b) (4, 0, 5)

In Exercises 7–10, find the coordinates of the point.

- 7. The point is located three units behind the *yz*-plane, four units to the right of the *xz*-plane, and five units above the *xy*-plane.
- **8.** The point is located seven units in front of the *yz*-plane, two units to the left of the *xz*-plane, and one unit below the *xy*-plane.
- **9.** The point is located on the *x*-axis, 12 units in front of the *yz*-plane.
- **10.** The point is located in the *yz*-plane, three units to the right of the *xz*-plane, and two units above the *xy*-plane.
- **11.** *Think About It* What is the *z*-coordinate of any point in the *xy*-plane?
- **12.** *Think About It* What is the *x*-coordinate of any point in the *yz*-plane?

In Exercises 13–24, determine the location of a point (x, y, z) that satisfies the condition(s).

13. $z = 6$	14. $y = 2$
15. $x = -3$	16. $z = -\frac{5}{2}$
17. $y < 0$	18. $x > 0$
19. $ y \leq 3$	20. $ x > 4$
21. $xy > 0, z = -3$	22. $xy < 0, z = 4$
23. $xyz < 0$	24. $xyz > 0$

In Exercises 25–28, find the distance between the points.

25. (0, 0, 0), (-4, 2, 7) **26.** (-2, 3, 2), (2, -5, -2) **27.** (1, -2, 4), (6, -2, -2)**28.** (2, 2, 3), (4, -5, 6) In Exercises 29–32, find the lengths of the sides of the triangle with the indicated vertices, and determine whether the triangle is a right triangle, an isosceles triangle, or neither.

- **29.** (0, 0, 4), (2, 6, 7), (6, 4, -8)
- **30.** (3, 4, 1), (0, 6, 2), (3, 5, 6)
- **31.** (-1, 0, -2), (-1, 5, 2), (-3, -1, 1)
- **32.** (4, -1, -1), (2, 0, -4), (3, 5, -1)
- **33.** *Think About It* The triangle in Exercise 29 is translated five units upward along the *z*-axis. Determine the coordinates of the translated triangle.
- **34.** *Think About It* The triangle in Exercise 30 is translated three units to the right along the *y*-axis. Determine the coordinates of the translated triangle.

In Exercises 35 and 36, find the coordinates of the midpoint of the line segment joining the points.

35. $(5, -9, 7), (-2, 3, 3)$ 36. (4)	4, 0, -6), (8, 8, 20)
--	-----------------------

In Exercises 37-40, find the standard equation of the sphere.

37. Center: (0, 2, 5)	38. Center: (4, -1, 1)
Radius: 2	Radius: 5

- **39.** Endpoints of a diameter: (2, 0, 0), (0, 6, 0)
- **40.** Center: (-3, 2, 4), tangent to the *yz*-plane

In Exercises 41–44, complete the square to write the equation of the sphere in standard form. Find the center and radius.

41. $x^2 + y^2 + z^2 - 2x + 6y + 8z + 1 = 0$ **42.** $x^2 + y^2 + z^2 + 9x - 2y + 10z + 19 = 0$ **43.** $9x^2 + 9y^2 + 9z^2 - 6x + 18y + 1 = 0$ **44.** $4x^2 + 4y^2 + 4z^2 - 24x - 4y + 8z - 23 = 0$

In Exercises 45–48, describe the solid satisfying the condition.

45.
$$x^2 + y^2 + z^2 \le 36$$

46. $x^2 + y^2 + z^2 > 4$
47. $x^2 + y^2 + z^2 < 4x - 6y + 8z - 13$
48. $x^2 + y^2 + z^2 > -4x + 6y - 8z - 13$

In Exercises 49–52, (a) find the component form of the vector v, (b) write the vector using standard unit vector notation, and (c) sketch the vector with its initial point at the origin.





In Exercises 53-56, find the component form and magnitude of the vector v with the given initial and terminal points. Then find a unit vector in the direction of v.

Initial Point	Terminal Poin
53. (3, 2, 0)	(4, 1, 6)
54. (4, -5, 2)	(-1, 7, -3)
55. (-4, 3, 1)	(-5, 3, 0)
56. (1, −2, 4)	(2, 4, -2)

In Exercises 57 and 58, the initial and terminal points of a vector v are given. (a) Sketch the directed line segment, (b) find the component form of the vector, (c) write the vector using standard unit vector notation, and (d) sketch the vector with its initial point at the origin.

57. Initial point: $(-1, 2, 3)$	58. Initial point: $(2, -1, -2)$
Terminal point: (3, 3, 4)	Terminal point: $(-4, 3, 7)$

In Exercises 59 and 60, the vector v and its initial point are given. Find the terminal point.

59. $\mathbf{v} = \langle 3, -5, 6 \rangle$ Initial point: (0, 6, 2) **60.** $\mathbf{v} = \langle 1, -\frac{2}{3}, \frac{1}{2} \rangle$ Initial point: (0, 2, $\frac{5}{2}$)

In Exercises 61 and 62, find each scalar multiple of v and sketch its graph.

61. $\mathbf{v} = \langle 1, 2, 2 \rangle$		62. $\mathbf{v} = \langle 2, -$	$\langle 2, -2, 1 \rangle$	
(a) 2 v	(b) – v	(a) –v	(b) 2 v	
(c) $\frac{3}{2}$ v	(d) 0 v	(c) $\frac{1}{2}$ v	(d) $\frac{5}{2}$ v	

In Exercises 63–68, find the vector z, given that $u = \langle 1, 2, 3 \rangle$, $v = \langle 2, 2, -1 \rangle$, and $w = \langle 4, 0, -4 \rangle$.

63. $z = u - v$	64. $z = u - v + 2w$
65. $z = 2u + 4v - w$	66. $z = 5u - 3v - \frac{1}{2}w$
67. $2z - 3u = w$	68. $2u + v - w + 3z = 0$

In Exercises 69–72, determine which of the vectors is (are) parallel to z. Use a graphing utility to confirm your results.

69. $z = \langle 3, 2, -5 \rangle$	70. $\mathbf{z} = \frac{1}{2}\mathbf{i} - \frac{2}{3}\mathbf{j} + \frac{3}{4}\mathbf{k}$
(a) $\langle -6, -4, 10 \rangle$	(a) $6i - 4j + 9k$
(b) $\left< 2, \frac{4}{3}, -\frac{10}{3} \right>$	(b) $-i + \frac{4}{3}j - \frac{3}{2}k$
(c) $\langle 6, 4, 10 \rangle$	(c) $12i + 9k$
(d) $(1, -4, 2)$	(d) $\frac{3}{4}i - j + \frac{9}{8}k$

- **71. z** has initial point (1, -1, 3) and terminal point (-2, 3, 5). (a) $-6\mathbf{i} + 8\mathbf{j} + 4\mathbf{k}$ (b) $4\mathbf{j} + 2\mathbf{k}$
- 72. z has initial point (5, 4, 1) and terminal point (-2, -4, 4).
 (a) (7, 6, 2)
 (b) (14, 16, -6)

In Exercises 73–76, use vectors to determine whether the points are collinear.

73. (0, -2, -5), (3, 4, 4), (2, 2, 1)
74. (4, -2, 7), (-2, 0, 3), (7, -3, 9)
75. (1, 2, 4), (2, 5, 0), (0, 1, 5)
76. (0, 0, 0), (1, 3, -2), (2, -6, 4)

In Exercises 77 and 78, use vectors to show that the points form the vertices of a parallelogram.

77. (2, 9, 1), (3, 11, 4), (0, 10, 2), (1, 12, 5) **78.** (1, 1, 3), (9, -1, -2), (11, 2, -9), (3, 4, -4)

In Exercises 79-84, find the magnitude of v.

79. $\mathbf{v} = \langle 0, 0, 0 \rangle$	80. $\mathbf{v} = \langle 1, 0, 3 \rangle$
81. $v = 3j - 5k$	82. $v = 2i + 5j - k$
83. $v = i - 2i - 3k$	84. $v = -4i + 3j + 7k$

In Exercises 85-88, find a unit vector (a) in the direction of v and (b) in the direction opposite of v.

85.	$\mathbf{v} = \langle 2, -1, 2 \rangle$	86.	$\mathbf{v} = \langle$	6, 0,	$8\rangle$
87.	$\mathbf{v} = \langle 3, 2, -5 \rangle$	88.	$\mathbf{v} = \langle$	8, 0,	$0\rangle$

89. *Programming* You are given the component forms of the vectors **u** and **v**. Write a program for a graphing utility in which the output is (a) the component form of $\mathbf{u} + \mathbf{v}$, (b) $\|\mathbf{u} + \mathbf{v}\|$, (c) $\|\mathbf{u}\|$, and (d) $\|\mathbf{v}\|$. (e) Run the program for the vectors $\mathbf{u} = \langle -1, 3, 4 \rangle$ and $\mathbf{v} = \langle 5, 4.5, -6 \rangle$.

CAPSTONE

90. Consider the two nonzero vectors **u** and **v**, and let *s* and *t* be real numbers. Describe the geometric figure generated by the terminal points of the three vectors $t\mathbf{v}$, $\mathbf{u} + t\mathbf{v}$, and $s\mathbf{u} + t\mathbf{v}$.

In Exercises 91 and 92, determine the values of *c* that satisfy the equation. Let u = -i + 2j + 3k and v = 2i + 2j - k.

91.
$$||c\mathbf{v}|| = 7$$
 92. $||c\mathbf{u}|| = 4$

In Exercises 93–96, find the vector v with the given magnitude and direction u.

	Magnitude	Direction
93.	10	$\mathbf{u} = \langle 0, 3, 3 \rangle$
94.	3	$\mathbf{u} = \langle 1, 1, 1 \rangle$
95.	$\frac{3}{2}$	$\mathbf{u} = \langle 2, -2, 1 \rangle$
96.	7	$\mathbf{u} = \langle -4, 6, 2 \rangle$

Licensed to:

782 Chapter 11 Vectors and the Geometry of Space

In Exercises 97 and 98, sketch the vector **v** and write its component form.

- **97.** v lies in the *yz*-plane, has magnitude 2, and makes an angle of 30° with the positive *y*-axis.
- **98.** v lies in the *xz*-plane, has magnitude 5, and makes an angle of 45° with the positive *z*-axis.

In Exercises 99 and 100, use vectors to find the point that lies two-thirds of the way from P to Q.

- **99.** P(4, 3, 0), Q(1, -3, 3) **100.** P(1, 2, 5), Q(6, 8, 2)
- 101. Let $\mathbf{u} = \mathbf{i} + \mathbf{j}$, $\mathbf{v} = \mathbf{j} + \mathbf{k}$, and $\mathbf{w} = a\mathbf{u} + b\mathbf{v}$.
 - (a) Sketch **u** and **v**.
 - (b) If $\mathbf{w} = \mathbf{0}$, show that *a* and *b* must both be zero.
 - (c) Find *a* and *b* such that $\mathbf{w} = \mathbf{i} + 2\mathbf{j} + \mathbf{k}$.
 - (d) Show that no choice of a and b yields $\mathbf{w} = \mathbf{i} + 2\mathbf{j} + 3\mathbf{k}$.
- **102.** *Writing* The initial and terminal points of the vector **v** are (x_1, y_1, z_1) and (x, y, z). Describe the set of all points (x, y, z) such that $\|\mathbf{v}\| = 4$.

WRITING ABOUT CONCEPTS

- **103.** A point in the three-dimensional coordinate system has coordinates (x_0, y_0, z_0) . Describe what each coordinate measures.
- **104.** Give the formula for the distance between the points (x_1, y_1, z_1) and (x_2, y_2, z_2) .
- **105.** Give the standard equation of a sphere of radius r, centered at (x_0, y_0, z_0) .
- 106. State the definition of parallel vectors.
- 107. Let A, B, and C be vertices of a triangle. Find $\overrightarrow{AB} + \overrightarrow{BC} + \overrightarrow{CA}$.
- **108.** Let $\mathbf{r} = \langle x, y, z \rangle$ and $\mathbf{r}_0 = \langle 1, 1, 1 \rangle$. Describe the set of all points (x, y, z) such that $\|\mathbf{r} \mathbf{r}_0\| = 2$.
- **109.** Numerical, Graphical, and Analytic Analysis The lights in an auditorium are 24-pound discs of radius 18 inches. Each disc is supported by three equally spaced cables that are L inches long (see figure).



- (a) Write the tension *T* in each cable as a function of *L*. Determine the domain of the function.
- (b) Use a graphing utility and the function in part (a) to complete the table.

L	20	25	30	35	40	45	50
T							

- (c) Use a graphing utility to graph the function in part (a). Determine the asymptotes of the graph.
- (d) Confirm the asymptotes of the graph in part (c) analytically.
- (e) Determine the minimum length of each cable if a cable is designed to carry a maximum load of 10 pounds.
- **110.** *Think About It* Suppose the length of each cable in Exercise 109 has a fixed length L = a, and the radius of each disc is r_0 inches. Make a conjecture about the limit $\lim_{r_0 \to a^-} T$ and give a reason for your answer.
- **111.** *Diagonal of a Cube* Find the component form of the unit vector **v** in the direction of the diagonal of the cube shown in the figure.



Figure for 111

Figure for 112

- **112.** *Tower Guy Wire* The guy wire supporting a 100-foot tower has a tension of 550 pounds. Using the distances shown in the figure, write the component form of the vector **F** representing the tension in the wire.
- **113.** *Load Supports* Find the tension in each of the supporting cables in the figure if the weight of the crate is 500 newtons.



Figure for 113

Figure for 114

- **114.** *Construction* A precast concrete wall is temporarily kept in its vertical position by ropes (see figure). Find the total force exerted on the pin at position *A*. The tensions in *AB* and *AC* are 420 pounds and 650 pounds.
- 115. Write an equation whose graph consists of the set of points P(x, y, z) that are twice as far from A(0, -1, 1) as from B(1, 2, 0).



- Use properties of the dot product of two vectors.
- Find the angle between two vectors using the dot product.
- Find the direction cosines of a vector in space.
- Find the projection of a vector onto another vector.
- Use vectors to find the work done by a constant force.

The Dot Product

So far you have studied two operations with vectors—vector addition and multiplication by a scalar—each of which yields another vector. In this section you will study a third vector operation, called the **dot product.** This product yields a scalar, rather than a vector.

DEFINITION OF DOT PRODUCT
The dot product of $\mathbf{u} = \langle u_1, u_2 \rangle$ and $\mathbf{v} = \langle v_1, v_2 \rangle$ is
$\mathbf{u} \cdot \mathbf{v} = u_1 v_1 + u_2 v_2.$
The dot product of $\mathbf{u} = \langle u_1, u_2, u_3 \rangle$ and $\mathbf{v} = \langle v_1, v_2, v_3 \rangle$ is
$\mathbf{u} \cdot \mathbf{v} = u_1 v_1 + u_2 v_2 + u_3 v_3.$

NOTE Because the dot product of two vectors yields a scalar, it is also called the **scalar product** (or **inner product**) of the two vectors.

THEOREM 11.4 PROPERTIES OF THE DOT PRODUCT

Let \mathbf{u} , \mathbf{v} , and \mathbf{w} be vectors in the plane or in space and let c be a scalar. **1.** $\mathbf{u} \cdot \mathbf{v} = \mathbf{v} \cdot \mathbf{u}$ Commutative Property **2.** $\mathbf{u} \cdot (\mathbf{v} + \mathbf{w}) = \mathbf{u} \cdot \mathbf{v} + \mathbf{u} \cdot \mathbf{w}$ Distributive Property **3.** $c(\mathbf{u} \cdot \mathbf{v}) = c\mathbf{u} \cdot \mathbf{v} = \mathbf{u} \cdot c\mathbf{v}$ **4.** $\mathbf{0} \cdot \mathbf{v} = 0$ **5.** $\mathbf{v} \cdot \mathbf{v} = \|\mathbf{v}\|^2$

PROOF To prove the first property, let $\mathbf{u} = \langle u_1, u_2, u_3 \rangle$ and $\mathbf{v} = \langle v_1, v_2, v_3 \rangle$. Then

$$\mathbf{u} \cdot \mathbf{v} = u_1 v_1 + u_2 v_2 + u_3 v_3$$

= $v_1 u_1 + v_2 u_2 + v_3 u_3$

For the fifth property, let $\mathbf{v} = \langle v_1, v_2, v_3 \rangle$. Then

$$\mathbf{v} \cdot \mathbf{v} = v_1^2 + v_2^2 + v_3^2 = \left(\sqrt{v_1^2 + v_2^2 + v_3^2}\right)^2 = \|\mathbf{v}\|^2.$$

Proofs of the other properties are left to you.

EXPLORATION

Interpreting a Dot Product Several vectors are shown below on the unit circle. Find the dot products of several pairs of vectors. Then find the angle between each pair that you used. Make a conjecture about the relationship between the dot product of two vectors and the angle between the vectors.



EXAMPLE 1 Finding Dot Products

Given $\mathbf{u} = \langle 2, -2 \rangle$, $\mathbf{v} = \langle 5, 8 \rangle$, and $\mathbf{w} = \langle -4, 3 \rangle$, find each of the following.

a. u · v	b. $(\mathbf{u} \cdot \mathbf{v})\mathbf{w}$	
c. u · (2 v)	d. $\ \mathbf{w}\ ^2$	
Solution		
a. $\mathbf{u} \cdot \mathbf{v} = \langle 2,$	$-2\rangle \cdot \langle 5, 8 \rangle = 2(5) + (-2)(8)$	= -6
b. $(\mathbf{u} \cdot \mathbf{v})\mathbf{w} =$	$-6\langle -4,3\rangle = \langle 24,-18\rangle$	
c. $\mathbf{u} \cdot (2\mathbf{v}) = 2$	$2(\mathbf{u} \cdot \mathbf{v}) = 2(-6) = -12$	Theorem 11.4
d. $\ \mathbf{w}\ ^2 = \mathbf{w} \cdot$	W	Theorem 11.4
$=\langle -4\rangle$	$4,3\rangle\cdot\langle-4,3\rangle$	Substitute $\langle -4, 3 \rangle$ for w .
=(-4)	(-4) + (3)(3)	Definition of dot product
= 25		Simplify.

Notice that the result of part (b) is a *vector* quantity, whereas the results of the other three parts are *scalar* quantities.

Angle Between Two Vectors

The **angle between two nonzero vectors** is the angle θ , $0 \le \theta \le \pi$, between their respective standard position vectors, as shown in Figure 11.24. The next theorem shows how to find this angle using the dot product. (Note that the angle between the zero vector and another vector is not defined here.)

THEOREM 11.5 ANGLE BETWEEN TWO VECTORS

If θ is the angle between two nonzero vectors **u** and **v**, then

$$\cos \theta = \frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{u}\| \|\mathbf{v}\|}$$

PROOF Consider the triangle determined by vectors \mathbf{u} , \mathbf{v} , and $\mathbf{v} - \mathbf{u}$, as shown in Figure 11.24. By the Law of Cosines, you can write

 $\|\mathbf{v} - \mathbf{u}\|^2 = \|\mathbf{u}\|^2 + \|\mathbf{v}\|^2 - 2\|\mathbf{u}\| \|\mathbf{v}\| \cos \theta.$

Using the properties of the dot product, the left side can be rewritten as

$$\|\mathbf{v} - \mathbf{u}\|^2 = (\mathbf{v} - \mathbf{u}) \cdot (\mathbf{v} - \mathbf{u})$$
$$= (\mathbf{v} - \mathbf{u}) \cdot \mathbf{v} - (\mathbf{v} - \mathbf{u}) \cdot \mathbf{u}$$
$$= \mathbf{v} \cdot \mathbf{v} - \mathbf{u} \cdot \mathbf{v} - \mathbf{v} \cdot \mathbf{u} + \mathbf{u} \cdot \mathbf{u}$$
$$= \|\mathbf{v}\|^2 - 2\mathbf{u} \cdot \mathbf{v} + \|\mathbf{u}\|^2$$

and substitution back into the Law of Cosines yields

$$\|\mathbf{v}\|^{2} - 2\mathbf{u} \cdot \mathbf{v} + \|\mathbf{u}\|^{2} = \|\mathbf{u}\|^{2} + \|\mathbf{v}\|^{2} - 2\|\mathbf{u}\| \|\mathbf{v}\| \cos \theta$$
$$-2\mathbf{u} \cdot \mathbf{v} = -2\|\mathbf{u}\| \|\mathbf{v}\| \cos \theta$$
$$\cos \theta = \frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{u}\| \|\mathbf{v}\|}.$$



If the angle between two vectors is known, rewriting Theorem 11.5 in the form

 $\mathbf{u} \cdot \mathbf{v} = \|\mathbf{u}\| \|\mathbf{v}\| \cos \theta$

Alternative form of dot product

produces an alternative way to calculate the dot product. From this form, you can see that because $\|\mathbf{u}\|$ and $\|\mathbf{v}\|$ are always positive, $\mathbf{u} \cdot \mathbf{v}$ and $\cos \theta$ will always have the same sign. Figure 11.25 shows the possible orientations of two vectors.



From Theorem 11.5, you can see that two nonzero vectors meet at a right angle if and only if their dot product is zero. Two such vectors are said to be **orthogonal**.

DEFINITION OF ORTHOGONAL VECTORS

The vectors **u** and **v** are orthogonal if $\mathbf{u} \cdot \mathbf{v} = 0$.

NOTE The terms "perpendicular," "orthogonal," and "normal" all mean essentially the same thing—meeting at right angles. However, it is common to say that two vectors are *orthogonal*, two lines or planes are *perpendicular*, and a vector is *normal* to a given line or plane.

lchap

From this definition, it follows that the zero vector is orthogonal to every vector **u**, because $\mathbf{0} \cdot \mathbf{u} = 0$. Moreover, for $0 \le \theta \le \pi$, you know that $\cos \theta = 0$ if and only if $\theta = \pi/2$. So, you can use Theorem 11.5 to conclude that two *nonzero* vectors are orthogonal if and only if the angle between them is $\pi/2$.

EXAMPLE 2 Finding the Angle Between Two Vectors

For $\mathbf{u} = \langle 3, -1, 2 \rangle$, $\mathbf{v} = \langle -4, 0, 2 \rangle$, $\mathbf{w} = \langle 1, -1, -2 \rangle$, and $\mathbf{z} = \langle 2, 0, -1 \rangle$, find the angle between each pair of vectors.

a.
$$\mathbf{u}$$
 and \mathbf{v} **b.** \mathbf{u} and \mathbf{w} **c.** \mathbf{v} and \mathbf{z}

Solution

a.
$$\cos \theta = \frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{u}\| \|\mathbf{v}\|} = \frac{-12 + 0 + 4}{\sqrt{14}\sqrt{20}} = \frac{-8}{2\sqrt{14}\sqrt{5}} = \frac{-4}{\sqrt{70}}$$

Because $\mathbf{u} \cdot \mathbf{v} < 0$, $\theta = \arccos \frac{4}{\sqrt{70}} \approx 2.069$ radians.

b. $\cos \theta = \frac{\mathbf{u} \cdot \mathbf{w}}{\|\mathbf{u}\| \|\mathbf{w}\|} = \frac{3+1-4}{\sqrt{14}\sqrt{6}} = \frac{0}{\sqrt{84}} = 0$

Because $\mathbf{u} \cdot \mathbf{w} = 0$, \mathbf{u} and \mathbf{w} are *orthogonal*. So, $\theta = \pi/2$.

c.
$$\cos \theta = \frac{\mathbf{v} \cdot \mathbf{z}}{\|\mathbf{v}\| \|\mathbf{z}\|} = \frac{-8 + 0 - 2}{\sqrt{20}\sqrt{5}} = \frac{-10}{\sqrt{100}} = -1$$

Consequently, $\theta = \pi$. Note that v and z are parallel, with v = -2z.



Direction angles Figure 11.26

Direction Cosines

For a vector in the plane, you have seen that it is convenient to measure direction in terms of the angle, measured counterclockwise, from the positive x-axis to the vector. In space it is more convenient to measure direction in terms of the angles between the nonzero vector **v** and the three unit vectors **i**, **j**, and **k**, as shown in Figure 11.26. The angles α , β , and γ are the **direction angles of v**, and $\cos \alpha$, $\cos \beta$, and $\cos \gamma$ are the direction cosines of v. Because

$$\mathbf{v} \cdot \mathbf{i} = \|\mathbf{v}\| \|\mathbf{i}\| \cos \alpha = \|\mathbf{v}\| \cos \alpha$$

and

$$\mathbf{v} \cdot \mathbf{i} = \langle v_1, v_2, v_3 \rangle \cdot \langle 1, 0, 0 \rangle = v$$

it follows that $\cos \alpha = v_1 / \|\mathbf{v}\|$. By similar reasoning with the unit vectors **j** and **k**, you have

$$\cos \alpha = \frac{v_1}{\|\mathbf{v}\|} \qquad \qquad \alpha \text{ is the angle between } \mathbf{v} \text{ and } \mathbf{i}.$$

$$\cos \beta = \frac{v_2}{\|\mathbf{v}\|} \qquad \qquad \beta \text{ is the angle between } \mathbf{v} \text{ and } \mathbf{j}.$$

$$\cos \gamma = \frac{v_3}{\|\mathbf{v}\|}.$$

$$\gamma \text{ is the angle between } \mathbf{v} \text{ and } \mathbf{k}.$$

Consequently, any nonzero vector \mathbf{v} in space has the normalized form

$$\frac{\mathbf{v}}{\|\mathbf{v}\|} = \frac{v_1}{\|\mathbf{v}\|}\mathbf{i} + \frac{v_2}{\|\mathbf{v}\|}\mathbf{j} + \frac{v_3}{\|\mathbf{v}\|}\mathbf{k} = \cos\alpha\mathbf{i} + \cos\beta\mathbf{j} + \cos\gamma\mathbf{k}$$

and because $\mathbf{v}/\|\mathbf{v}\|$ is a unit vector, it follows that

 $\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1.$

EXAMPLE 3 Finding Direction Angles

Find the direction cosines and angles for the vector $\mathbf{v} = 2\mathbf{i} + 3\mathbf{j} + 4\mathbf{k}$, and show that $\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1.$

Solution Because $\|\mathbf{v}\| = \sqrt{2^2 + 3^2 + 4^2} = \sqrt{29}$, you can write the following.

$$\cos \alpha = \frac{v_1}{\|\mathbf{v}\|} = \frac{2}{\sqrt{29}} \implies \alpha \approx 68.2^{\circ}$$
Angle between **v** and **i**
$$\cos \beta = \frac{v_2}{\|\mathbf{v}\|} = \frac{3}{\sqrt{29}} \implies \beta \approx 56.1^{\circ}$$
Angle between **v** and **j**
$$\cos \gamma = \frac{v_3}{\|\mathbf{v}\|} = \frac{4}{\sqrt{29}} \implies \gamma \approx 42.0^{\circ}$$
Angle between **v** and **k**

Furthermore, the sum of the squares of the direction cosines is

$$\cos^{2} \alpha + \cos^{2} \beta + \cos^{2} \gamma = \frac{4}{29} + \frac{9}{29} + \frac{16}{29}$$
$$= \frac{29}{29}$$
$$= 1.$$

The direction angles of v **Figure 11.27**

Copyright 2010 Cengage Learning, Inc. All Rights Reserved. May not be copied, scanned, or duplicated, in whole or in part.



See Figure 11.27.



The force due to gravity pulls the boat against the ramp and down the ramp. **Figure 11.28**

Projections and Vector Components

You have already seen applications in which two vectors are added to produce a resultant vector. Many applications in physics and engineering pose the reverse problem—decomposing a given vector into the sum of two **vector components.** The following physical example enables you to see the usefulness of this procedure.

Consider a boat on an inclined ramp, as shown in Figure 11.28. The force **F** due to gravity pulls the boat *down* the ramp and *against* the ramp. These two forces, \mathbf{w}_1 and \mathbf{w}_2 , are orthogonal—they are called the vector components of **F**.

 $\mathbf{F} = \mathbf{w}_1 + \mathbf{w}_2$ Vector components of \mathbf{F}

The forces \mathbf{w}_1 and \mathbf{w}_2 help you analyze the effect of gravity on the boat. For example, \mathbf{w}_1 indicates the force necessary to keep the boat from rolling down the ramp, whereas \mathbf{w}_2 indicates the force that the tires must withstand.

DEFINITIONS OF PROJECTION AND VECTOR COMPONENTS

Let **u** and **v** be nonzero vectors. Moreover, let $\mathbf{u} = \mathbf{w}_1 + \mathbf{w}_2$, where \mathbf{w}_1 is parallel to **v**, and \mathbf{w}_2 is orthogonal to **v**, as shown in Figure 11.29.

- w₁ is called the projection of u onto v or the vector component of u along v, and is denoted by w₁ = proj_vu.
- 2. $\mathbf{w}_2 = \mathbf{u} \mathbf{w}_1$ is called the vector component of u orthogonal to v.



 $w_1 = \text{proj}_v u = \text{projection of } u \text{ onto } v = \text{vector component of } u \text{ along } v$ $w_2 = \text{vector component of } u \text{ orthogonal to } v$ Figure 11.29





Find the vector component of $\mathbf{u} = \langle 5, 10 \rangle$ that is orthogonal to $\mathbf{v} = \langle 4, 3 \rangle$, given that $\mathbf{w}_1 = \text{proj}_{\mathbf{v}} \mathbf{u} = \langle 8, 6 \rangle$ and

$$\mathbf{u} = \langle 5, 10 \rangle = \mathbf{w}_1 + \mathbf{w}_2$$

Solution Because $\mathbf{u} = \mathbf{w}_1 + \mathbf{w}_2$, where \mathbf{w}_1 is parallel to \mathbf{v} , it follows that \mathbf{w}_2 is the vector component of \mathbf{u} orthogonal to \mathbf{v} . So, you have

$$\mathbf{w}_2 = \mathbf{u} - \mathbf{w}_1$$

= $\langle 5, 10 \rangle - \langle 8, 6 \rangle$
= $\langle -3, 4 \rangle$.

Check to see that \mathbf{w}_2 is orthogonal to \mathbf{v} , as shown in Figure 11.30.

 $u = w_1 + w_2$ Figure 11.30

788

NOTE Note the distinction between the terms "component" and "vector component." For example, using the standard unit vectors with $\mathbf{u} = u_1 \mathbf{i} + u_2 \mathbf{j}$, u_1 is the *component* of \mathbf{u} in the direction of \mathbf{i} and $u_1 \mathbf{i}$ is the *vector component* in the direction of \mathbf{i} .







From Example 4, you can see that it is easy to find the vector component \mathbf{w}_2 once you have found the projection, \mathbf{w}_1 , of **u** onto **v**. To find this projection, use the dot product given in the theorem below, which you will prove in Exercise 92.

THEOREM 11.6 PROJECTION USING THE DOT PRODUCT

If \mathbf{u} and \mathbf{v} are nonzero vectors, then the projection of \mathbf{u} onto \mathbf{v} is given by

 $\operatorname{proj}_{\mathbf{v}}\mathbf{u} = \left(\frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{v}\|^2}\right)\mathbf{v}.$

The projection of \mathbf{u} onto \mathbf{v} can be written as a scalar multiple of a unit vector in the direction of \mathbf{v} . That is,

$$\left(\frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{v}\|^2}\right)\mathbf{v} = \left(\frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{v}\|}\right)\frac{\mathbf{v}}{\|\mathbf{v}\|} = (k)\frac{\mathbf{v}}{\|\mathbf{v}\|} \implies k = \frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{v}\|} = \|\mathbf{u}\|\cos\theta.$$

The scalar *k* is called the **component of u in the direction of v.**

EXAMPLE 5 Decomposing a Vector into Vector Components

Find the projection of **u** onto **v** and the vector component of **u** orthogonal to **v** for the vectors $\mathbf{u} = 3\mathbf{i} - 5\mathbf{j} + 2\mathbf{k}$ and $\mathbf{v} = 7\mathbf{i} + \mathbf{j} - 2\mathbf{k}$ shown in Figure 11.31.

Solution The projection of **u** onto **v** is

$$\mathbf{w}_1 = \left(\frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{v}\|^2}\right) \mathbf{v} = \left(\frac{12}{54}\right) (7\mathbf{i} + \mathbf{j} - 2\mathbf{k}) = \frac{14}{9}\mathbf{i} + \frac{2}{9}\mathbf{j} - \frac{4}{9}\mathbf{k}.$$

The vector component of \mathbf{u} orthogonal to \mathbf{v} is the vector

$$\mathbf{w}_2 = \mathbf{u} - \mathbf{w}_1 = (3\mathbf{i} - 5\mathbf{j} + 2\mathbf{k}) - (\frac{14}{9}\mathbf{i} + \frac{2}{9}\mathbf{j} - \frac{4}{9}\mathbf{k}) = \frac{13}{9}\mathbf{i} - \frac{47}{9}\mathbf{j} + \frac{22}{9}\mathbf{k}$$

EXAMPLE 6 Finding a Force

A 600-pound boat sits on a ramp inclined at 30°, as shown in Figure 11.32. What force is required to keep the boat from rolling down the ramp?

Solution Because the force due to gravity is vertical and downward, you can represent the gravitational force by the vector $\mathbf{F} = -600\mathbf{j}$. To find the force required to keep the boat from rolling down the ramp, project \mathbf{F} onto a unit vector \mathbf{v} in the direction of the ramp, as follows.

$$\mathbf{v} = \cos 30^{\circ}\mathbf{i} + \sin 30^{\circ}\mathbf{j} = \frac{\sqrt{3}}{2}\mathbf{i} + \frac{1}{2}\mathbf{j}$$
 Unit vector along ramp

Therefore, the projection of \mathbf{F} onto \mathbf{v} is given by

$$\mathbf{w}_1 = \operatorname{proj}_{\mathbf{v}} \mathbf{F} = \left(\frac{\mathbf{F} \cdot \mathbf{v}}{\|\mathbf{v}\|^2}\right) \mathbf{v} = (\mathbf{F} \cdot \mathbf{v}) \mathbf{v} = (-600) \left(\frac{1}{2}\right) \mathbf{v} = -300 \left(\frac{\sqrt{3}}{2}\mathbf{i} + \frac{1}{2}\mathbf{j}\right).$$

The magnitude of this force is 300, and therefore a force of 300 pounds is required to keep the boat from rolling down the ramp.



Work = $||\mathbf{F}|| || \overrightarrow{PQ} ||$ (a) Force acts along the line of motion.



Work = $\| \operatorname{proj}_{\overline{PQ}} \mathbf{F} \| \| \overline{PQ} \|$

(b) Force acts at angle θ with the line of motion. **Figure 11.33**



Figure 11.34



See www.CalcChat.com for worked-out solutions to odd-numbered exercises.

In Exercises 1–8, find (a) $u\cdot v,$ (b) $u\cdot u,$ (c) $\|u\|^2,$ (d) $(u\cdot v)v,$ and (e) $u\cdot$ (2v).

1. $\mathbf{u} = \langle 3, 4 \rangle$, $\mathbf{v} = \langle -1, 5 \rangle$ **2.** $\mathbf{u} = \langle 4, 10 \rangle$, $\mathbf{v} = \langle -2, 3 \rangle$ **3.** $\mathbf{u} = \langle 6, -4 \rangle$, $\mathbf{v} = \langle -3, 2 \rangle$ **4.** $\mathbf{u} = \langle -4, 8 \rangle$, $\mathbf{v} = \langle 7, 5 \rangle$ **5.** $\mathbf{u} = \langle 2, -3, 4 \rangle$, $\mathbf{v} = \langle 0, 6, 5 \rangle$ **6.** $\mathbf{u} = \mathbf{i}$, $\mathbf{v} = \mathbf{i}$ **7.** $\mathbf{u} = 2\mathbf{i} - \mathbf{j} + \mathbf{k}$ $\mathbf{v} = \mathbf{i} - \mathbf{k}$ **8.** $\mathbf{u} = 2\mathbf{i} + \mathbf{j} - 2\mathbf{k}$ $\mathbf{v} = \mathbf{i} - 3\mathbf{j} + 2\mathbf{k}$

In Exercises 9 and 10, find u $\,\cdot\,$ v.

9. $\|\mathbf{u}\| = 8$, $\|\mathbf{v}\| = 5$, and the angle between \mathbf{u} and \mathbf{v} is $\pi/3$. 10. $\|\mathbf{u}\| = 40$, $\|\mathbf{v}\| = 25$, and the angle between \mathbf{u} and \mathbf{v} is $5\pi/6$.

In Exercises 11–18, find the angle θ between the vectors.

11. $\mathbf{u} = \langle 1, 1 \rangle, \mathbf{v} = \langle 2, -2 \rangle$ **12.** $\mathbf{u} = \langle 3, 1 \rangle, \mathbf{v} = \langle 2, -1 \rangle$

Work

The work W done by the constant force \mathbf{F} acting along the line of motion of an object is given by

 $W = (\text{magnitude of force})(\text{distance}) = \|\mathbf{F}\| \| \overline{PQ} \|$

as shown in Figure 11.33(a). If the constant force \mathbf{F} is not directed along the line of motion, you can see from Figure 11.33(b) that the work *W* done by the force is

$$W = \|\operatorname{proj}_{\overrightarrow{PQ}}\mathbf{F}\| \| \overrightarrow{PQ} \| = (\cos \theta) \|\mathbf{F}\| \| \overrightarrow{PQ} \| = \mathbf{F} \cdot \overrightarrow{PQ}.$$

This notion of work is summarized in the following definition.

DEFINITION OF WORKThe work W done by a constant force F as its point of application moves
along the vector \overrightarrow{PQ} is given by either of the following.1. $W = \| \operatorname{proj}_{\overrightarrow{PQ}} F \| \| \overrightarrow{PQ} \|$ Projection form2. $W = F \cdot \overrightarrow{PQ}$ Dot product form

EXAMPLE 7 Finding Work

To close a sliding door, a person pulls on a rope with a constant force of 50 pounds at a constant angle of 60° , as shown in Figure 11.34. Find the work done in moving the door 12 feet to its closed position.

Solution Using a projection, you can calculate the work as follows.



13. $\mathbf{u} = 3\mathbf{i} + \mathbf{j}, \mathbf{v} = -2\mathbf{i} + 4\mathbf{j}$ 14. $\mathbf{u} = \cos\left(\frac{\pi}{6}\right)\mathbf{i} + \sin\left(\frac{\pi}{6}\right)\mathbf{j}$ $\mathbf{v} = \cos\left(\frac{3\pi}{4}\right)\mathbf{i} + \sin\left(\frac{3\pi}{4}\right)\mathbf{j}$ 15. $\mathbf{u} = \langle 1, 1, 1 \rangle$ $\mathbf{v} = \langle 2, 1, -1 \rangle$ 16. $\mathbf{u} = 3\mathbf{i} + 2\mathbf{j} + \mathbf{k}$ $\mathbf{v} = 2\mathbf{i} - 3\mathbf{j}$ 17. $\mathbf{u} = 3\mathbf{i} + 4\mathbf{j}$ $\mathbf{v} = -2\mathbf{j} + 3\mathbf{k}$ 18. $\mathbf{u} = 2\mathbf{i} - 2\mathbf{j} + \mathbf{k}$

In Exercises 19–26, determine whether u and v are orthogonal, parallel, or neither.

19.
$$\mathbf{u} = \langle 4, 0 \rangle, \quad \mathbf{v} = \langle 1, 1 \rangle$$

20. $\mathbf{u} = \langle 2, 18 \rangle, \quad \mathbf{v} = \left\langle \frac{3}{2}, -\frac{1}{6} \right\rangle$

21.
$$\mathbf{u} = \langle 4, 3 \rangle$$

 $\mathbf{v} = \left\langle \frac{1}{2}, -\frac{2}{3} \right\rangle$ **22.** $\mathbf{u} = -\frac{1}{3}(\mathbf{i} - 2\mathbf{j})$
 $\mathbf{v} = 2\mathbf{i} - 4\mathbf{j}$ **23.** $\mathbf{u} = \mathbf{j} + 6\mathbf{k}$
 $\mathbf{v} = \mathbf{i} - 2\mathbf{j} - \mathbf{k}$ **24.** $\mathbf{u} = -2\mathbf{i} + 3\mathbf{j} - \mathbf{k}$
 $\mathbf{v} = 2\mathbf{i} + \mathbf{j} - \mathbf{k}$ **25.** $\mathbf{u} = \langle 2, -3, 1 \rangle$
 $\mathbf{v} = \langle -1, -1, -1 \rangle$ **26.** $\mathbf{u} = \langle \cos \theta, \sin \theta, -1 \rangle$
 $\mathbf{v} = \langle \sin \theta, -\cos \theta, 0 \rangle$

In Exercises 27–30, the vertices of a triangle are given. Determine whether the triangle is an acute triangle, an obtuse triangle, or a right triangle. Explain your reasoning.

```
27. (1, 2, 0), (0, 0, 0), (-2, 1, 0)
28. (-3, 0, 0), (0, 0, 0), (1, 2, 3)
29. (2, 0, 1), (0, 1, 2), (-0.5, 1.5, 0)
30. (2, -7, 3), (-1, 5, 8), (4, 6, -1)
```

In Exercises 31–34, find the direction cosines of u and demonstrate that the sum of the squares of the direction cosines is 1.

31. $\mathbf{u} = \mathbf{i} + 2\mathbf{j} + 2\mathbf{k}$ **32.** $\mathbf{u} = 5\mathbf{i} + 3\mathbf{j} - \mathbf{k}$ **33.** $\mathbf{u} = \langle 0, 6, -4 \rangle$ **34.** $\mathbf{u} = \langle a, b, c \rangle$

In Exercises 35–38, find the direction angles of the vector.

35.	$\mathbf{u} = 3\mathbf{i} + 2\mathbf{j} - 2\mathbf{k}$	36. $\mathbf{u} = -4\mathbf{i} + 3\mathbf{j} + 5\mathbf{k}$
37.	$\mathbf{u} = \langle -1, 5, 2 \rangle$	38. $\mathbf{u} = \langle -2, 6, 1 \rangle$

In Exercises 39 and 40, use a graphing utility to find the magnitude and direction angles of the resultant of forces F_1 and F_2 with initial points at the origin. The magnitude and terminal point of each vector are given.

Vector	Magnitude	Terminal Point
39. F ₁	50 lb	(10, 5, 3)
\mathbf{F}_2	80 lb	(12, 7, -5)
40. F ₁	300 N	(-20, -10, 5)
\mathbf{F}_2	100 N	(5, 15, 0)

41. *Load-Supporting Cables* A load is supported by three cables, as shown in the figure. Find the direction angles of the load-supporting cable *OA*.



42. *Load-Supporting Cables* The tension in the cable *OA* in Exercise 41 is 200 newtons. Determine the weight of the load.

In Exercises 43–50, (a) find the projection of u onto v, and (b) find the vector component of u orthogonal to v.

43. $\mathbf{u} = \langle 6, 7 \rangle$, $\mathbf{v} = \langle 1, 4 \rangle$ 44. $\mathbf{u} = \langle 9, 7 \rangle$, $\mathbf{v} = \langle 1, 3 \rangle$ 45. $\mathbf{u} = 2\mathbf{i} + 3\mathbf{j}$, $\mathbf{v} = 5\mathbf{i} + \mathbf{j}$ 46. $\mathbf{u} = 2\mathbf{i} - 3\mathbf{j}$, $\mathbf{v} = 3\mathbf{i} + 2\mathbf{j}$ 47. $\mathbf{u} = \langle 0, 3, 3 \rangle$, $\mathbf{v} = \langle -1, 1, 1 \rangle$ 48. $\mathbf{u} = \langle 8, 2, 0 \rangle$, $\mathbf{v} = \langle 2, 1, -1 \rangle$ 49. $\mathbf{u} = 2\mathbf{i} + \mathbf{j} + 2\mathbf{k}$, $\mathbf{v} = 3\mathbf{j} + 4\mathbf{k}$ 50. $\mathbf{u} = \mathbf{i} + 4\mathbf{k}$, $\mathbf{v} = 3\mathbf{i} + 2\mathbf{k}$

WRITING ABOUT CONCEPTS

- **51.** Define the dot product of vectors \mathbf{u} and \mathbf{v} .
- **52.** State the definition of orthogonal vectors. If vectors are neither parallel nor orthogonal, how do you find the angle between them? Explain.
- **53.** Determine which of the following are defined for nonzero vectors **u**, **v**, and **w**. Explain your reasoning.

(a)
$$\mathbf{u} \cdot (\mathbf{v} + \mathbf{w})$$
 (b) $(\mathbf{u} \cdot \mathbf{v})\mathbf{w}$

- (c) $\mathbf{u} \cdot \mathbf{v} + \mathbf{w}$ (d) $\|\mathbf{u}\| \cdot (\mathbf{v} + \mathbf{w})$
- 54. Describe direction cosines and direction angles of a vector \mathbf{v} .
- **55.** Give a geometric description of the projection of \mathbf{u} onto \mathbf{v} .
- 56. What can be said about the vectors u and v if (a) the projection of u onto v equals u and (b) the projection of u onto v equals 0?
- 57. If the projection of u onto v has the same magnitude as the projection of v onto u, can you conclude that ||u|| = ||v||? Explain.

CAPSTONE

58. What is known about θ , the angle between two nonzero vectors **u** and **v**, if

(a) $\mathbf{u} \cdot \mathbf{v} = 0$? (b) $\mathbf{u} \cdot \mathbf{v} > 0$? (c) $\mathbf{u} \cdot \mathbf{v} < 0$?

- **59.** *Revenue* The vector $\mathbf{u} = \langle 3240, 1450, 2235 \rangle$ gives the numbers of hamburgers, chicken sandwiches, and cheeseburgers, respectively, sold at a fast-food restaurant in one week. The vector $\mathbf{v} = \langle 1.35, 2.65, 1.85 \rangle$ gives the prices (in dollars) per unit for the three food items. Find the dot product $\mathbf{u} \cdot \mathbf{v}$, and explain what information it gives.
- **60.** *Revenue* Repeat Exercise 59 after increasing prices by 4%. Identify the vector operation used to increase prices by 4%.
- 61. Programming Given vectors u and v in component form, write a program for a graphing utility in which the output is (a) ||u||, (b) ||v||, and (c) the angle between u and v.
- **62.** *Programming* Use the program you wrote in Exercise 61 to find the angle between the vectors $\mathbf{u} = \langle 8, -4, 2 \rangle$ and $\mathbf{v} = \langle 2, 5, 2 \rangle$.

- **63.** *Programming* Given vectors **u** and **v** in component form, write a program for a graphing utility in which the output is the component form of the projection of **u** onto **v**.
- **64.** *Programming* Use the program you wrote in Exercise 63 to find the projection of **u** onto **v** for $\mathbf{u} = \langle 5, 6, 2 \rangle$ and $\mathbf{v} = \langle -1, 3, 4 \rangle$.

Think About It In Exercises 65 and 66, use the figure to determine mentally the projection of u onto v. (The coordinates of the terminal points of the vectors in standard position are given.) Verify your results analytically.



In Exercises 67–70, find two vectors in opposite directions that are orthogonal to the vector u. (The answers are not unique.)

67.	$\mathbf{u} = -\frac{1}{4}\mathbf{i} + \frac{3}{2}\mathbf{j}$	68. $u = 9i - 4j$
69.	$\mathbf{u} = \langle 3, 1, -2 \rangle$	70. $\mathbf{u} = \langle 4, -3, 6 \rangle$

71. *Braking Load* A 48,000-pound truck is parked on a 10° slope (see figure). Assume the only force to overcome is that due to gravity. Find (a) the force required to keep the truck from rolling down the hill and (b) the force perpendicular to the hill.





Figure for 72

- **72.** *Load-Supporting Cables* Find the magnitude of the projection of the load-supporting cable *OA* onto the positive *z*-axis as shown in the figure.
- **73.** *Work* An object is pulled 10 feet across a floor, using a force of 85 pounds. The direction of the force is 60° above the horizontal (see figure). Find the work done.





Figure for 73

Figure for 74

- **74.** *Work* A toy wagon is pulled by exerting a force of 25 pounds on a handle that makes a 20° angle with the horizontal (see figure in left column). Find the work done in pulling the wagon 50 feet.
- **75.** *Work* A car is towed using a force of 1600 newtons. The chain used to pull the car makes a 25° angle with the horizontal. Find the work done in towing the car 2 kilometers.
- **76.** *Work* A sled is pulled by exerting a force of 100 newtons on a rope that makes a 25° angle with the horizontal. Find the work done in pulling the sled 40 meters.

True or False? In Exercises 77 and 78, determine whether the statement is true or false. If it is false, explain why or give an example that shows it is false.

- 77. If $\mathbf{u} \cdot \mathbf{v} = \mathbf{u} \cdot \mathbf{w}$ and $\mathbf{u} \neq \mathbf{0}$, then $\mathbf{v} = \mathbf{w}$.
- **78.** If **u** and **v** are orthogonal to **w**, then $\mathbf{u} + \mathbf{v}$ is orthogonal to **w**.
- **79.** Find the angle between a cube's diagonal and one of its edges.
- **80.** Find the angle between the diagonal of a cube and the diagonal of one of its sides.

In Exercises 81–84, (a) find all points of intersection of the graphs of the two equations, (b) find the unit tangent vectors to each curve at their points of intersection, and (c) find the angles $(0^{\circ} \le \theta \le 90^{\circ})$ between the curves at their points of intersection.

- 81. $y = x^2$, $y = x^{1/3}$ 82. $y = x^3$, $y = x^{1/3}$ 83. $y = 1 - x^2$, $y = x^2 - 1$ 84. $(y + 1)^2 = x$, $y = x^3 - 1$
- **85.** Use vectors to prove that the diagonals of a rhombus are perpendicular.
- **86.** Use vectors to prove that a parallelogram is a rectangle if and only if its diagonals are equal in length.
- **87.** *Bond Angle* Consider a regular tetrahedron with vertices (0, 0, 0), (k, k, 0), (k, 0, k), and (0, k, k), where k is a positive real number.
 - (a) Sketch the graph of the tetrahedron.
 - (b) Find the length of each edge.
 - (c) Find the angle between any two edges.
 - (d) Find the angle between the line segments from the centroid (k/2, k/2, k/2) to two vertices. This is the bond angle for a molecule such as CH₄ or PbCl₄, where the structure of the molecule is a tetrahedron.
- **88.** Consider the vectors $\mathbf{u} = \langle \cos \alpha, \sin \alpha, 0 \rangle$ and $\mathbf{v} = \langle \cos \beta, \sin \beta, 0 \rangle$, where $\alpha > \beta$. Find the dot product of the vectors and use the result to prove the identity

 $\cos(\alpha - \beta) = \cos \alpha \cos \beta + \sin \alpha \sin \beta.$

- **89.** Prove that $\|\mathbf{u} \mathbf{v}\|^2 = \|\mathbf{u}\|^2 + \|\mathbf{v}\|^2 2\mathbf{u} \cdot \mathbf{v}$.
- **90.** Prove the Cauchy-Schwarz Inequality $|\mathbf{u} \cdot \mathbf{v}| \le ||\mathbf{u}|| ||\mathbf{v}||$.
- 91. Prove the triangle inequality $\|\mathbf{u} + \mathbf{v}\| \le \|\mathbf{u}\| + \|\mathbf{v}\|$.
- 92. Prove Theorem 11.6.

11.4 The Cross Product of Two Vectors in Space

- Find the cross product of two vectors in space.
- Use the triple scalar product of three vectors in space.

The Cross Product

EXPLORATION

Geometric Property of the Cross Product Three pairs of vectors are shown below. Use the definition to find the cross product of each pair. Sketch all three vectors in a three-dimensional system. Describe any relationships among the three vectors. Use your description to write a conjecture about \mathbf{u} , \mathbf{v} , and $\mathbf{u} \times \mathbf{v}$.







c. $\mathbf{u} = \langle 3, 3, 0 \rangle, \ \mathbf{v} = \langle 3, -3, 0 \rangle$



Many applications in physics, engineering, and geometry involve finding a vector in space that is orthogonal to two given vectors. In this section you will study a product that will yield such a vector. It is called the **cross product**, and it is most conveniently defined and calculated using the standard unit vector form. Because the cross product yields a vector, it is also called the **vector product**.

DEFINITION OF CROSS PRODUCT OF TWO VECTORS IN SPACE

Let

$$\mathbf{u} = u_1\mathbf{i} + u_2\mathbf{j} + u_3\mathbf{k}$$
 and $\mathbf{v} = v_1\mathbf{i} + v_2\mathbf{j} + v_3\mathbf{k}$

be vectors in space. The **cross product** of **u** and **v** is the vector

$$\mathbf{u} \times \mathbf{v} = (u_2 v_3 - u_3 v_2)\mathbf{i} - (u_1 v_3 - u_3 v_1)\mathbf{j} + (u_1 v_2 - u_2 v_1)\mathbf{k}.$$

NOTE Be sure you see that this definition applies only to three-dimensional vectors. The cross product is not defined for two-dimensional vectors.

A convenient way to calculate $\mathbf{u} \times \mathbf{v}$ is to use the following *determinant form* with cofactor expansion. (This 3 × 3 determinant form is used simply to help remember the formula for the cross product—it is technically not a determinant because the entries of the corresponding matrix are not all real numbers.)

$$\mathbf{u} \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ u_1 & u_2 & u_3 \\ v_1 & v_2 & v_3 \end{vmatrix} \longrightarrow \operatorname{Put} "u" \text{ in Row 2.} \\ \longrightarrow \operatorname{Put} "v" \text{ in Row 3.} \\ = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ u_1 & u_2 & u_3 \\ v_1 & v_2 & v_3 \end{vmatrix} \mathbf{i} - \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ u_1 & u_2 & u_3 \\ v_1 & v_2 & v_3 \end{vmatrix} \mathbf{j} + \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ u_1 & u_2 & u_3 \\ v_1 & v_2 & v_3 \end{vmatrix} \mathbf{k} \\ = \begin{vmatrix} u_2 & u_3 \\ v_2 & v_3 \end{vmatrix} \mathbf{i} - \begin{vmatrix} u_1 & u_3 \\ v_1 & v_3 \end{vmatrix} \mathbf{j} + \begin{vmatrix} u_1 & u_2 \\ v_1 & v_2 \end{vmatrix} \mathbf{k} \\ = (u_2 v_3 - u_3 v_2) \mathbf{i} - (u_1 v_3 - u_3 v_1) \mathbf{j} + (u_1 v_2 - u_2 v_1) \mathbf{k} \end{vmatrix}$$

Note the minus sign in front of the **j**-component. Each of the three 2×2 determinants can be evaluated by using the following diagonal pattern.

$$\begin{vmatrix} a & b \\ c & d \end{vmatrix} = ad - bc$$

Here are a couple of examples.

$$\begin{vmatrix} 2 & 4 \\ 3 & -1 \end{vmatrix} = (2)(-1) - (4)(3) = -2 - 12 = -14$$
$$\begin{vmatrix} 4 & 0 \\ -6 & 3 \end{vmatrix} = (4)(3) - (0)(-6) = 12$$

NOTATION FOR DOT AND CROSS PRODUCTS

The notation for the dot product and cross product of vectors was first introduced by the American physicist Josiah Willard Gibbs (1839–1903). In the early 1880s, Gibbs built a system to represent physical quantities called "vector analysis." The system was a departure from Hamilton's theory of quaternions.

EXAMPLE 1 Finding the Cross Product

b. $\mathbf{v} \times \mathbf{u}$

Given $\mathbf{u} = \mathbf{i} - 2\mathbf{j} + \mathbf{k}$ and $\mathbf{v} = 3\mathbf{i} + \mathbf{j} - 2\mathbf{k}$, find each of the following.

c. $\mathbf{v} \times \mathbf{v}$

a. $\mathbf{u} \times \mathbf{v}$ Solution

a. $\mathbf{u} \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & -2 & 1 \\ 3 & 1 & -2 \end{vmatrix} = \begin{vmatrix} -2 & 1 \\ 1 & -2 \end{vmatrix} \mathbf{i} - \begin{vmatrix} 1 & 1 \\ 3 & -2 \end{vmatrix} \mathbf{j} + \begin{vmatrix} 1 & -2 \\ 3 & 1 \end{vmatrix} \mathbf{k}$ $= (4 - 1)\mathbf{i} - (-2 - 3)\mathbf{j} + (1 + 6)\mathbf{k}$ $= 3\mathbf{i} + 5\mathbf{j} + 7\mathbf{k}$ **b.** $\mathbf{v} \times \mathbf{u} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 3 & 1 & -2 \\ 1 & -2 & 1 \end{vmatrix} = \begin{vmatrix} 1 & -2 \\ -2 & 1 \end{vmatrix} \mathbf{i} - \begin{vmatrix} 3 & -2 \\ 1 & 1 \end{vmatrix} \mathbf{j} + \begin{vmatrix} 3 & 1 \\ 1 & -2 \end{vmatrix} \mathbf{k}$ $= (1 - 4)\mathbf{i} - (3 + 2)\mathbf{j} + (-6 - 1)\mathbf{k}$ $= -3\mathbf{i} - 5\mathbf{j} - 7\mathbf{k}$

Note that this result is the negative of that in part (a).

	i	j	k	
c. $\mathbf{v} \times \mathbf{v} =$	3	1	-2	= 0
	3	1	-2	

The results obtained in Example 1 suggest some interesting *algebraic* properties of the cross product. For instance, $\mathbf{u} \times \mathbf{v} = -(\mathbf{v} \times \mathbf{u})$, and $\mathbf{v} \times \mathbf{v} = \mathbf{0}$. These properties, and several others, are summarized in the following theorem.

THEOREM 11.7 ALGEBRAIC PROPERTIES OF THE CROSS PRODUCT

Let **u**, **v**, and **w** be vectors in space, and let *c* be a scalar.

1. $\mathbf{u} \times \mathbf{v} = -(\mathbf{v} \times \mathbf{u})$ 2. $\mathbf{u} \times (\mathbf{v} + \mathbf{w}) = (\mathbf{u} \times \mathbf{v}) + (\mathbf{u} \times \mathbf{w})$ 3. $c(\mathbf{u} \times \mathbf{v}) = (c\mathbf{u}) \times \mathbf{v} = \mathbf{u} \times (c\mathbf{v})$ 4. $\mathbf{u} \times \mathbf{0} = \mathbf{0} \times \mathbf{u} = \mathbf{0}$ 5. $\mathbf{u} \times \mathbf{u} = \mathbf{0}$ 6. $\mathbf{u} \cdot (\mathbf{v} \times \mathbf{w}) = (\mathbf{u} \times \mathbf{v}) \cdot \mathbf{w}$

PROOF To prove Property 1, let $\mathbf{u} = u_1\mathbf{i} + u_2\mathbf{j} + u_3\mathbf{k}$ and $\mathbf{v} = v_1\mathbf{i} + v_2\mathbf{j} + v_3\mathbf{k}$. Then,

$$\mathbf{u} \times \mathbf{v} = (u_2 v_3 - u_3 v_2)\mathbf{i} - (u_1 v_3 - u_3 v_1)\mathbf{j} + (u_1 v_2 - u_2 v_1)\mathbf{k}$$

and

$$\mathbf{v} \times \mathbf{u} = (v_2 u_3 - v_3 u_2)\mathbf{i} - (v_1 u_3 - v_3 u_1)\mathbf{j} + (v_1 u_2 - v_2 u_1)\mathbf{k}$$

which implies that $\mathbf{u} \times \mathbf{v} = -(\mathbf{v} \times \mathbf{u})$. Proofs of Properties 2, 3, 5, and 6 are left as exercises (see Exercises 59–62).

794 Chapter 11 Vectors and the Geometry of Space

NOTE It follows from Properties 1 and 2 in Theorem 11.8 that if **n** is a unit vector orthogonal to both **u** and **v**, then

 $\mathbf{u} \times \mathbf{v} = \pm (\|\mathbf{u}\| \|\mathbf{v}\| \sin \theta) \mathbf{n}.$

Note that Property 1 of Theorem 11.7 indicates that the cross product is *not* commutative. In particular, this property indicates that the vectors $\mathbf{u} \times \mathbf{v}$ and $\mathbf{v} \times \mathbf{u}$ have equal lengths but opposite directions. The following theorem lists some other geometric properties of the cross product of two vectors.

THEOREM 11.8 GEOMETRIC PROPERTIES OF THE CROSS PRODUCT

Let **u** and **v** be nonzero vectors in space, and let θ be the angle between **u** and **v**.

- **1.** $\mathbf{u} \times \mathbf{v}$ is orthogonal to both \mathbf{u} and \mathbf{v} .
- **2.** $\|\mathbf{u} \times \mathbf{v}\| = \|\mathbf{u}\| \|\mathbf{v}\| \sin \theta$
- 3. $\mathbf{u} \times \mathbf{v} = \mathbf{0}$ if and only if \mathbf{u} and \mathbf{v} are scalar multiples of each other.
- **4.** $\|\mathbf{u} \times \mathbf{v}\|$ = area of parallelogram having \mathbf{u} and \mathbf{v} as adjacent sides.

PROOF To prove Property 2, note because $\cos \theta = (\mathbf{u} \cdot \mathbf{v})/(||\mathbf{u}|| ||\mathbf{v}||)$, it follows that

$$\begin{aligned} \|\mathbf{u}\| \|\mathbf{v}\| \sin \theta &= \|\mathbf{u}\| \|\mathbf{v}\| \sqrt{1 - \cos^2 \theta} \\ &= \|\mathbf{u}\| \|\mathbf{v}\| \sqrt{1 - \frac{(\mathbf{u} \cdot \mathbf{v})^2}{\|\mathbf{u}\|^2 \|\mathbf{v}\|^2}} \\ &= \sqrt{\|\mathbf{u}\|^2 \|\mathbf{v}\|^2 - (\mathbf{u} \cdot \mathbf{v})^2} \\ &= \sqrt{(u_1^2 + u_2^2 + u_3^2)(v_1^2 + v_2^2 + v_3^2) - (u_1v_1 + u_2v_2 + u_3v_3)^2} \\ &= \sqrt{(u_2v_3 - u_3v_2)^2 + (u_1v_3 - u_3v_1)^2 + (u_1v_2 - u_2v_1)^2} \\ &= \|\mathbf{u} \times \mathbf{v}\|. \end{aligned}$$

To prove Property 4, refer to Figure 11.35, which is a parallelogram having **v** and **u** as adjacent sides. Because the height of the parallelogram is $\|\mathbf{v}\| \sin \theta$, the area is

Area = (base)(height)
=
$$\|\mathbf{u}\| \|\mathbf{v}\| \sin \theta$$

= $\|\mathbf{u} \times \mathbf{v}\|$.

Proofs of Properties 1 and 3 are left as exercises (see Exercises 63 and 64).

Both $\mathbf{u} \times \mathbf{v}$ and $\mathbf{v} \times \mathbf{u}$ are perpendicular to the plane determined by \mathbf{u} and \mathbf{v} . One way to remember the orientations of the vectors \mathbf{u} , \mathbf{v} , and $\mathbf{u} \times \mathbf{v}$ is to compare them with the unit vectors \mathbf{i} , \mathbf{j} , and $\mathbf{k} = \mathbf{i} \times \mathbf{j}$, as shown in Figure 11.36. The three vectors \mathbf{u} , \mathbf{v} , and $\mathbf{u} \times \mathbf{v}$ form a *right-handed system*, whereas the three vectors \mathbf{u} , \mathbf{v} , and $\mathbf{v} \times \mathbf{u}$ form a *left-handed system*.



Right-handed systems **Figure 11.36**



The vectors **u** and **v** form adjacent sides of a parallelogram. Figure 11.35

n



The vector $\mathbf{u} \times \mathbf{v}$ is orthogonal to both \mathbf{u} and \mathbf{v} . Figure 11.37



The area of the parallelogram is approximately 32.19. Figure 11.38

EXAMPLE 2 Using the Cross Product

Find a unit vector that is orthogonal to both

 $\mathbf{u} = \mathbf{i} - 4\mathbf{j} + \mathbf{k}$ and $\mathbf{v} = 2\mathbf{i} + 3\mathbf{j}$.

Solution The cross product $\mathbf{u} \times \mathbf{v}$, as shown in Figure 11.37, is orthogonal to both \mathbf{u} and \mathbf{v} .

$$\mathbf{u} \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & -4 & 1 \\ 2 & 3 & 0 \end{vmatrix}$$

= $-3\mathbf{i} + 2\mathbf{j} + 11\mathbf{k}$ Cross product

Because

$$\|\mathbf{u} \times \mathbf{v}\| = \sqrt{(-3)^2 + 2^2 + 11^2} = \sqrt{134}$$

a unit vector orthogonal to both **u** and **v** is

$$\frac{\mathbf{u} \times \mathbf{v}}{\|\mathbf{u} \times \mathbf{v}\|} = -\frac{3}{\sqrt{134}}\mathbf{i} + \frac{2}{\sqrt{134}}\mathbf{j} + \frac{11}{\sqrt{134}}\mathbf{k}.$$

NOTE In Example 2, note that you could have used the cross product $\mathbf{v} \times \mathbf{u}$ to form a unit vector that is orthogonal to both \mathbf{u} and \mathbf{v} . With that choice, you would have obtained the negative of the unit vector found in the example.

EXAMPLE 3 Geometric Application of the Cross Product

Show that the quadrilateral with vertices at the following points is a parallelogram, and find its area.

$$A = (5, 2, 0) \qquad B = (2, 6, 1) C = (2, 4, 7) \qquad D = (5, 0, 6)$$

Solution From Figure 11.38 you can see that the sides of the quadrilateral correspond to the following four vectors.

$$\overrightarrow{AB} = -3\mathbf{i} + 4\mathbf{j} + \mathbf{k} \qquad \overrightarrow{CD} = 3\mathbf{i} - 4\mathbf{j} - \mathbf{k} = -\overrightarrow{AB}$$

$$\overrightarrow{AD} = 0\mathbf{i} - 2\mathbf{j} + 6\mathbf{k} \qquad \overrightarrow{CB} = 0\mathbf{i} + 2\mathbf{j} - 6\mathbf{k} = -\overrightarrow{AD}$$

So, \overrightarrow{AB} is parallel to \overrightarrow{CD} and \overrightarrow{AD} is parallel to \overrightarrow{CB} , and you can conclude that the quadrilateral is a parallelogram with \overrightarrow{AB} and \overrightarrow{AD} as adjacent sides. Moreover, because

$$\overrightarrow{AB} \times \overrightarrow{AD} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -3 & 4 & 1 \\ 0 & -2 & 6 \end{vmatrix}$$

$$= 26\mathbf{i} + 18\mathbf{j} + 6\mathbf{k}$$
Cross product

the area of the parallelogram is

$$\|\overline{AB} \times \overline{AD}\| = \sqrt{1036} \approx 32.19$$

Is the parallelogram a rectangle? You can determine whether it is by finding the angle between the vectors \overrightarrow{AB} and \overrightarrow{AD} .



The moment of **F** about *P* **Figure 11.39**



A vertical force of 50 pounds is applied at point Q. Figure 11.40

FOR FURTHER INFORMATION To see how the cross product is used to model the torque of the robot arm of a space shuttle, see the article "The Long Arm of Calculus" by Ethan Berkove and Rich Marchand in *The College Mathematics Journal*. To view this article, go to the website *www.matharticles.com*.

In physics, the cross product can be used to measure **torque**—the **moment M of a force F about a point** P**,** as shown in Figure 11.39. If the point of application of the force is Q, the moment of **F** about P is given by

$$\mathbf{M} = \overrightarrow{PQ} \times \mathbf{F}.$$

Moment of **F** about P

The magnitude of the moment **M** measures the tendency of the vector \overrightarrow{PQ} to rotate counterclockwise (using the right-hand rule) about an axis directed along the vector **M**.

EXAMPLE 4 An Application of the Cross Product

A vertical force of 50 pounds is applied to the end of a one-foot lever that is attached to an axle at point *P*, as shown in Figure 11.40. Find the moment of this force about the point *P* when $\theta = 60^{\circ}$.

Solution If you represent the 50-pound force as $\mathbf{F} = -50\mathbf{k}$ and the lever as

$$\overrightarrow{PQ} = \cos(60^\circ)\mathbf{j} + \sin(60^\circ)\mathbf{k} = \frac{1}{2}\mathbf{j} + \frac{\sqrt{3}}{2}\mathbf{k}$$

the moment of \mathbf{F} about *P* is given by

$$\mathbf{M} = \overrightarrow{PQ} \times \mathbf{F} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 0 & \frac{1}{2} & \frac{\sqrt{3}}{2} \\ 0 & 0 & -50 \end{vmatrix} = -25\mathbf{i}.$$
 Moment of **F** about *P*

The magnitude of this moment is 25 foot-pounds.

NOTE In Example 4, note that the moment (the tendency of the lever to rotate about its axle) is dependent on the angle θ . When $\theta = \pi/2$, the moment is 0. The moment is greatest when $\theta = 0$.

The Triple Scalar Product

For vectors **u**, **v**, and **w** in space, the dot product of **u** and $\mathbf{v} \times \mathbf{w}$

 $\mathbf{u} \cdot (\mathbf{v} \times \mathbf{w})$

is called the **triple scalar product**, as defined in Theorem 11.9. The proof of this theorem is left as an exercise (see Exercise 67).

THEOREM 11.9 THE TRIPLE SCALAR PRODUCT

For $\mathbf{u} = u_1 \mathbf{i} + u_2 \mathbf{j} + u_3 \mathbf{k}$, $\mathbf{v} = v_1 \mathbf{i} + v_2 \mathbf{j} + v_3 \mathbf{k}$, and $\mathbf{w} = w_1 \mathbf{i} + w_2 \mathbf{j} + w_3 \mathbf{k}$, the triple scalar product is given by

$$\mathbf{u} \cdot (\mathbf{v} \times \mathbf{w}) = \begin{vmatrix} u_1 & u_2 & u_3 \\ v_1 & v_2 & v_3 \\ w_1 & w_2 & w_3 \end{vmatrix}$$

NOTE The value of a determinant is multiplied by -1 if two rows are interchanged. After two such interchanges, the value of the determinant will be unchanged. So, the following triple scalar products are equivalent.

$$\mathbf{u} \cdot (\mathbf{v} \times \mathbf{w}) = \mathbf{v} \cdot (\mathbf{w} \times \mathbf{u}) = \mathbf{w} \cdot (\mathbf{u} \times \mathbf{v})$$


Area of base = $\|\mathbf{v} \times \mathbf{w}\|$ Volume of parallelepiped = $|\mathbf{u} \cdot (\mathbf{v} \times \mathbf{w})|$ Figure 11.41



The parallelepiped has a volume of 36. **Figure 11.42**

If the vectors \mathbf{u} , \mathbf{v} , and \mathbf{w} do not lie in the same plane, the triple scalar product $\mathbf{u} \cdot (\mathbf{v} \times \mathbf{w})$ can be used to determine the volume of the parallelepiped (a polyhedron, all of whose faces are parallelograms) with \mathbf{u} , \mathbf{v} , and \mathbf{w} as adjacent edges, as shown in Figure 11.41. This is established in the following theorem.

THEOREM 11.10 GEOMETRIC PROPERTY OF THE TRIPLE SCALAR PRODUCT

The volume V of a parallelepiped with vectors \mathbf{u} , \mathbf{v} , and \mathbf{w} as adjacent edges is given by

 $V = |\mathbf{u} \cdot (\mathbf{v} \times \mathbf{w})|.$

(PROOF) In Figure 11.41, note that

 $\|\mathbf{v} \times \mathbf{w}\| =$ area of base

and

 $\|\operatorname{proj}_{\mathbf{v}\times\mathbf{w}}\mathbf{u}\| = \text{height of parallelepiped.}$

Therefore, the volume is

$$V = (\text{height})(\text{area of base}) = \|\text{proj}_{\mathbf{v}\times\mathbf{w}}\mathbf{u}\|\|\mathbf{v}\times\mathbf{w}\|$$
$$= \left|\frac{\mathbf{u}\cdot(\mathbf{v}\times\mathbf{w})}{\|\mathbf{v}\times\mathbf{w}\|}\right|\|\mathbf{v}\times\mathbf{w}\|$$
$$= |\mathbf{u}\cdot(\mathbf{v}\times\mathbf{w})|.$$

EXAMPLE 5 Volume by the Triple Scalar Product

Find the volume of the parallelepiped shown in Figure 11.42 having $\mathbf{u} = 3\mathbf{i} - 5\mathbf{j} + \mathbf{k}$, $\mathbf{v} = 2\mathbf{j} - 2\mathbf{k}$, and $\mathbf{w} = 3\mathbf{i} + \mathbf{j} + \mathbf{k}$ as adjacent edges.

Solution By Theorem 11.10, you have

$$V = |\mathbf{u} \cdot (\mathbf{v} \times \mathbf{w})|$$
 Triple scalar product

$$= \begin{vmatrix} 3 & -5 & 1 \\ 0 & 2 & -2 \\ 3 & 1 & 1 \end{vmatrix}$$

$$= 3\begin{vmatrix} 2 & -2 \\ 1 & 1 \end{vmatrix} - (-5)\begin{vmatrix} 0 & -2 \\ 3 & 1 \end{vmatrix} + (1)\begin{vmatrix} 0 & 2 \\ 3 & 1 \end{vmatrix}$$

$$= 3(4) + 5(6) + 1(-6)$$

$$= 36.$$

A natural consequence of Theorem 11.10 is that the volume of the parallelepiped is 0 if and only if the three vectors are coplanar. That is, if the vectors $\mathbf{u} = \langle u_1, u_2, u_3 \rangle$, $\mathbf{v} = \langle v_1, v_2, v_3 \rangle$, and $\mathbf{w} = \langle w_1, w_2, w_3 \rangle$ have the same initial point, they lie in the same plane if and only if

$$\mathbf{u} \cdot (\mathbf{v} \times \mathbf{w}) = \begin{vmatrix} u_1 & u_2 & u_3 \\ v_1 & v_2 & v_3 \\ w_1 & w_2 & w_3 \end{vmatrix} = 0.$$

11.4 Exercises

See www.CalcChat.com for worked-out solutions to odd-numbered exercises.

In Exercises 1–6, find the cross product of the unit vectors and sketch your result.

1. j × i	2. i × j
3. $\mathbf{j} \times \mathbf{k}$	4. $\mathbf{k} \times \mathbf{j}$
5. i × k	6. $\mathbf{k} \times \mathbf{i}$

In Exercises 7–10, find (a) $u \times v$, (b) $v \times u$, and (c) $v \times v$.

7. $\mathbf{u} = -2\mathbf{i} + 4\mathbf{j}$	8. $u = 3i + 5k$
$\mathbf{v} = 3\mathbf{i} + 2\mathbf{j} + 5\mathbf{k}$	$\mathbf{v} = 2\mathbf{i} + 3\mathbf{j} - 2\mathbf{k}$
9. $\mathbf{u} = \langle 7, 3, 2 \rangle$	10. $\mathbf{u} = \langle 3, -2, -2 \rangle$
$\mathbf{v} = \langle 1, -1, 5 \rangle$	$\mathbf{v} = \langle 1, 5, 1 \rangle$

In Exercises 11–16, find $\mathbf{u}\times\mathbf{v}$ and show that it is orthogonal to both \mathbf{u} and $\mathbf{v}.$

11. $\mathbf{u} = \langle 12, -3, 0 \rangle$	12. $\mathbf{u} = \langle -1, 1, 2 \rangle$
$\mathbf{v} = \langle -2, 5, 0 \rangle$	$\mathbf{v} = \langle 0, 1, 0 \rangle$
13. $\mathbf{u} = \langle 2, -3, 1 \rangle$	14. $\mathbf{u} = \langle -10, 0, 6 \rangle$
$\mathbf{v} = \langle 1, -2, 1 \rangle$	$\mathbf{v} = \langle 5, -3, 0 \rangle$
15. $u = i + j + k$	16. $u = i + 6j$
$\mathbf{v} = 2\mathbf{i} + \mathbf{j} - \mathbf{k}$	$\mathbf{v} = -2\mathbf{i} + \mathbf{j} + \mathbf{k}$

Think About It In Exercises 17–20, use the vectors u and v shown in the figure to sketch a vector in the direction of the indicated cross product in a right-handed system.



17.	$\mathbf{u} \times \mathbf{v}$	18.	v	×	u
19.	$(-\mathbf{v}) imes \mathbf{u}$	20.	u	\times	$(\mathbf{u}\times\mathbf{v})$

CAS In Exercises 21–24, use a computer algebra system to find u × v and a unit vector orthogonal to u and v.

21. $\mathbf{u} = \langle 4, -3.5, 7 \rangle$	22. $\mathbf{u} = \langle -8, -6, 4 \rangle$
$\mathbf{v} = \langle 2.5, 9, 3 \rangle$	$\mathbf{v} = \langle 10, -12, -2 \rangle$
23. $\mathbf{u} = -3\mathbf{i} + 2\mathbf{j} - 5\mathbf{k}$	24. $u = 0.7k$
$\mathbf{v} = 0.4\mathbf{i} - 0.8\mathbf{j} + 0.2\mathbf{k}$	$\mathbf{v} = 1.5\mathbf{i} + 6.2\mathbf{k}$

- **25.** *Programming* Given the vectors **u** and **v** in component form, write a program for a graphing utility in which the output is $\mathbf{u} \times \mathbf{v}$ and $\|\mathbf{u} \times \mathbf{v}\|$.
- **26.** *Programming* Use the program you wrote in Exercise 25 to find $\mathbf{u} \times \mathbf{v}$ and $||\mathbf{u} \times \mathbf{v}||$ for $\mathbf{u} = \langle -2, 6, 10 \rangle$ and $\mathbf{v} = \langle 3, 8, 5 \rangle$.

Area In Exercises 27–30, find the area of the parallelogram that has the given vectors as adjacent sides. Use a computer algebra system or a graphing utility to verify your result.

27. u = j	28. $u = i + j + k$
$\mathbf{v} = \mathbf{j} + \mathbf{k}$	$\mathbf{v} = \mathbf{j} + \mathbf{k}$
29. $\mathbf{u} = \langle 3, 2, -1 \rangle$	30. $\mathbf{u} = \langle 2, -1, 0 \rangle$
$\mathbf{v} = \langle 1, 2, 3 \rangle$	$\mathbf{v} = \langle -1, 2, 0 \rangle$

Area In Exercises 31 and 32, verify that the points are the vertices of a parallelogram, and find its area.

31. *A*(0, 3, 2), *B*(1, 5, 5), *C*(6, 9, 5), *D*(5, 7, 2) **32.** *A*(2, -3, 1), *B*(6, 5, -1), *C*(7, 2, 2), *D*(3, -6, 4)

Area In Exercises 33–36, find the area of the triangle with the given vertices. (*Hint:* $\frac{1}{2} || \mathbf{u} \times \mathbf{v} ||$ is the area of the triangle having u and v as adjacent sides.)

- 33. A(0, 0, 0), B(1, 0, 3), C(-3, 2, 0)
 34. A(2, -3, 4), B(0, 1, 2), C(-1, 2, 0)
 35. A(2, -7, 3), B(-1, 5, 8), C(4, 6, -1)
 36. A(1, 2, 0), B(-2, 1, 0), C(0, 0, 0)
- **37.** *Torque* A child applies the brakes on a bicycle by applying a downward force of 20 pounds on the pedal when the crank makes a 40° angle with the horizontal (see figure). The crank is 6 inches in length. Find the torque at *P*.



Figure for 37

Figure for 38

- **38.** *Torque* Both the magnitude and the direction of the force on a crankshaft change as the crankshaft rotates. Find the torque on the crankshaft using the position and data shown in the figure.
- **39.** *Optimization* A force of 56 pounds acts on the pipe wrench shown in the figure on the next page.
 - (a) Find the magnitude of the moment about *O* by evaluating $\|\overrightarrow{OA} \times \mathbf{F}\|$. Use a graphing utility to graph the resulting function of θ .
 - (b) Use the result of part (a) to determine the magnitude of the moment when $\theta = 45^{\circ}$.
 - (c) Use the result of part (a) to determine the angle θ when the magnitude of the moment is maximum. Is the answer what you expected? Why or why not?



Figure for 39

- Figure for 40
- **40.** *Optimization* A force of 180 pounds acts on the bracket shown in the figure.
 - (a) Determine the vector \overrightarrow{AB} and the vector \mathbf{F} representing the force. (\mathbf{F} will be in terms of θ .)
 - (b) Find the magnitude of the moment about A by evaluating $\|\overline{AB} \times \mathbf{F}\|$.
 - (c) Use the result of part (b) to determine the magnitude of the moment when $\theta = 30^{\circ}$.
 - (d) Use the result of part (b) to determine the angle θ when the magnitude of the moment is maximum. At that angle, what is the relationship between the vectors **F** and \overrightarrow{AB} ? Is it what you expected? Why or why not?
- (e) Use a graphing utility to graph the function for the magnitude of the moment about A for $0^{\circ} \le \theta \le 180^{\circ}$. Find the zero of the function in the given domain. Interpret the meaning of the zero in the context of the problem.

In Exercises 41–44, find $u \cdot (v \times w)$.



Volume In Exercises 45 and 46, use the triple scalar product to find the volume of the parallelepiped having adjacent edges u, v, and w.



Volume In Exercises 47 and 48, find the volume of the parallelepiped with the given vertices.

- 47. (0, 0, 0), (3, 0, 0), (0, 5, 1), (2, 0, 5) (3, 5, 1), (5, 0, 5), (2, 5, 6), (5, 5, 6)
 48. (0, 0, 0), (0, 4, 0), (-3, 0, 0), (-1, 1, 5) (-3, 4, 0), (-1, 5, 5), (-4, 1, 5), (-4, 5, 5)
- **49.** If $\mathbf{u} \times \mathbf{v} = \mathbf{0}$ and $\mathbf{u} \cdot \mathbf{v} = \mathbf{0}$, what can you conclude about \mathbf{u} and \mathbf{v} ?
- **50.** Identify the dot products that are equal. Explain your reasoning. (Assume **u**, **v**, and **w** are nonzero vectors.)

WRITING ABOUT CONCEPTS

- **51.** Define the cross product of vectors **u** and **v**.
- **52.** State the geometric properties of the cross product.
- **53.** If the magnitudes of two vectors are doubled, how will the magnitude of the cross product of the vectors change? Explain.

CAPSTONE

54. The vertices of a triangle in space are (x_1, y_1, z_1) , (x_2, y_2, z_2) , and (x_3, y_3, z_3) . Explain how to find a vector perpendicular to the triangle.

True or False? In Exercises 55–58, determine whether the statement is true or false. If it is false, explain why or give an example that shows it is false.

- **55.** It is possible to find the cross product of two vectors in a two-dimensional coordinate system.
- 56. If u and v are vectors in space that are nonzero and nonparallel, then $\mathbf{u} \times \mathbf{v} = \mathbf{v} \times \mathbf{u}$.
- **57.** If $\mathbf{u} \neq \mathbf{0}$ and $\mathbf{u} \times \mathbf{v} = \mathbf{u} \times \mathbf{w}$, then $\mathbf{v} = \mathbf{w}$.

58. If $\mathbf{u} \neq \mathbf{0}$, $\mathbf{u} \cdot \mathbf{v} = \mathbf{u} \cdot \mathbf{w}$, and $\mathbf{u} \times \mathbf{v} = \mathbf{u} \times \mathbf{w}$, then $\mathbf{v} = \mathbf{w}$.

In Exercises 59–66, prove the property of the cross product.

59.
$$\mathbf{u} \times (\mathbf{v} + \mathbf{w}) = (\mathbf{u} \times \mathbf{v}) + (\mathbf{u} \times \mathbf{w})$$

60.
$$c(\mathbf{u} \times \mathbf{v}) = (c\mathbf{u}) \times \mathbf{v} = \mathbf{u} \times (c\mathbf{v})$$

- 61. $\mathbf{u} \times \mathbf{u} = \mathbf{0}$
- 62. $\mathbf{u} \cdot (\mathbf{v} \times \mathbf{w}) = (\mathbf{u} \times \mathbf{v}) \cdot \mathbf{w}$
- **63.** $\mathbf{u} \times \mathbf{v}$ is orthogonal to both \mathbf{u} and \mathbf{v} .
- **64.** $\mathbf{u} \times \mathbf{v} = \mathbf{0}$ if and only if \mathbf{u} and \mathbf{v} are scalar multiples of each other.
- **65.** Prove that $\|\mathbf{u} \times \mathbf{v}\| = \|\mathbf{u}\| \|\mathbf{v}\|$ if \mathbf{u} and \mathbf{v} are orthogonal.
- **66.** Prove that $\mathbf{u} \times (\mathbf{v} \times \mathbf{w}) = (\mathbf{u} \cdot \mathbf{w})\mathbf{v} (\mathbf{u} \cdot \mathbf{v})\mathbf{w}$.
- **67.** Prove Theorem 11.9.

11.5 Lines and Planes in Space



Line *L* and its direction vector v Figure 11.43



The vector **v** is parallel to the line *L*. **Figure 11.44**

- Write a set of parametric equations for a line in space.
- Write a linear equation to represent a plane in space.
- Sketch the plane given by a linear equation.
- Find the distances between points, planes, and lines in space.

Lines in Space

In the plane, *slope* is used to determine an equation of a line. In space, it is more convenient to use *vectors* to determine the equation of a line.

In Figure 11.43, consider the line *L* through the point $P(x_1, y_1, z_1)$ and parallel to the vector $\mathbf{v} = \langle a, b, c \rangle$. The vector \mathbf{v} is a **direction vector** for the line *L*, and *a*, *b*, and *c* are **direction numbers.** One way of describing the line *L* is to say that it consists of all points Q(x, y, z) for which the vector \overline{PQ} is parallel to \mathbf{v} . This means that \overline{PQ} is a scalar multiple of \mathbf{v} , and you can write $\overline{PQ} = t\mathbf{v}$, where *t* is a scalar (a real number).

$$\overline{PQ} = \langle x - x_1, y - y_1, z - z_1 \rangle = \langle at, bt, ct \rangle = t \mathbf{v}$$

By equating corresponding components, you can obtain **parametric equations** of a line in space.

THEOREM 11.11 PARAMETRIC EQUATIONS OF A LINE IN SPACE

A line *L* parallel to the vector $\mathbf{v} = \langle a, b, c \rangle$ and passing through the point $P(x_1, y_1, z_1)$ is represented by the **parametric equations**

 $x = x_1 + at$, $y = y_1 + bt$, and $z = z_1 + ct$.

If the direction numbers a, b, and c are all nonzero, you can eliminate the parameter t to obtain symmetric equations of the line.



EXAMPLE 1 Finding Parametric and Symmetric Equations

Find parametric and symmetric equations of the line *L* that passes through the point (1, -2, 4) and is parallel to $\mathbf{v} = \langle 2, 4, -4 \rangle$.

Solution To find a set of parametric equations of the line, use the coordinates $x_1 = 1$, $y_1 = -2$, and $z_1 = 4$ and direction numbers a = 2, b = 4, and c = -4 (see Figure 11.44).

$$x = 1 + 2t$$
, $y = -2 + 4t$, $z = 4 - 4t$ Parametric equations

Because a, b, and c are all nonzero, a set of symmetric equations is

$$\frac{x-1}{2} = \frac{y+2}{4} = \frac{z-4}{-4}.$$
 Symmetric equations

Neither parametric equations nor symmetric equations of a given line are unique. For instance, in Example 1, by letting t = 1 in the parametric equations you would obtain the point (3, 2, 0). Using this point with the direction numbers a = 2, b = 4, and c = -4 would produce a different set of parametric equations

x = 3 + 2t, y = 2 + 4t, and z = -4t.

EXAMPLE 2 Parametric Equations of a Line Through Two Points

Find a set of parametric equations of the line that passes through the points (-2, 1, 0) and (1, 3, 5).

Solution Begin by using the points P(-2, 1, 0) and Q(1, 3, 5) to find a direction vector for the line passing through P and Q, given by

$$\mathbf{v} = \overline{PQ} = \langle 1 - (-2), 3 - 1, 5 - 0 \rangle = \langle 3, 2, 5 \rangle = \langle a, b, c \rangle.$$

Using the direction numbers a = 3, b = 2, and c = 5 with the point P(-2, 1, 0), you can obtain the parametric equations

x = -2 + 3t, y = 1 + 2t, and z = 5t.

NOTE As *t* varies over all real numbers, the parametric equations in Example 2 determine the points (x, y, z) on the line. In particular, note that t = 0 and t = 1 give the original points (-2, 1, 0) and (1, 3, 5).

Planes in Space

You have seen how an equation of a line in space can be obtained from a point on the line and a vector *parallel* to it. You will now see that an equation of a plane in space can be obtained from a point in the plane and a vector *normal* (perpendicular) to the plane.

Consider the plane containing the point $P(x_1, y_1, z_1)$ having a nonzero normal vector $\mathbf{n} = \langle a, b, c \rangle$, as shown in Figure 11.45. This plane consists of all points Q(x, y, z) for which vector \overrightarrow{PQ} is orthogonal to \mathbf{n} . Using the dot product, you can write the following.

 $\mathbf{n} \cdot \overrightarrow{PQ} = 0$ $\langle a, b, c \rangle \cdot \langle x - x_1, y - y_1, z - z_1 \rangle = 0$ $a(x - x_1) + b(y - y_1) + c(z - z_1) = 0$

The third equation of the plane is said to be in standard form.

THEOREM 11.12 STANDARD EQUATION OF A PLANE IN SPACE

The plane containing the point (x_1, y_1, z_1) and having normal vector $\mathbf{n} = \langle a, b, c \rangle$ can be represented by the **standard form** of the equation of a plane

 $a(x - x_1) + b(y - y_1) + c(z - z_1) = 0.$

By regrouping terms, you obtain the general form of the equation of a plane in space.

ax + by + cz + d = 0

General form of equation of plane



The normal vector **n** is orthogonal to each vector \overrightarrow{PQ} in the plane. Figure 11.45



A plane determined by **u** and **v** Figure 11.46





The angle θ between two planes **Figure 11.47**

Given the general form of the equation of a plane, it is easy to find a normal vector to the plane. Simply use the coefficients of x, y, and z and write $\mathbf{n} = \langle a, b, c \rangle$.

EXAMPLE 3 Finding an Equation of a Plane in Three-Space

Find the general equation of the plane containing the points (2, 1, 1), (0, 4, 1), and (-2, 1, 4).

Solution To apply Theorem 11.12 you need a point in the plane and a vector that is normal to the plane. There are three choices for the point, but no normal vector is given. To obtain a normal vector, use the cross product of vectors \mathbf{u} and \mathbf{v} extending from the point (2, 1, 1) to the points (0, 4, 1) and (-2, 1, 4), as shown in Figure 11.46. The component forms of \mathbf{u} and \mathbf{v} are

$$\mathbf{u} = \langle 0 - 2, 4 - 1, 1 - 1 \rangle = \langle -2, 3, 0 \rangle$$
$$\mathbf{v} = \langle -2 - 2, 1 - 1, 4 - 1 \rangle = \langle -4, 0, 3 \rangle$$

and it follows that

$$\mathbf{n} = \mathbf{u} \times \mathbf{v}$$
$$= \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -2 & 3 & 0 \\ -4 & 0 & 3 \end{vmatrix}$$
$$= 9\mathbf{i} + 6\mathbf{j} + 12\mathbf{k}$$
$$= \langle a, b, c \rangle$$

is normal to the given plane. Using the direction numbers for **n** and the point $(x_1, y_1, z_1) = (2, 1, 1)$, you can determine an equation of the plane to be

$$a(x - x_1) + b(y - y_1) + c(z - z_1) = 0$$

$$9(x - 2) + 6(y - 1) + 12(z - 1) = 0$$

$$9x + 6y + 12z - 36 = 0$$

$$3x + 2y + 4z - 12 = 0.$$

Standard form
General form
Simplified general form

NOTE In Example 3, check to see that each of the three original points satisfies the equation 3x + 2y + 4z - 12 = 0.

Two distinct planes in three-space either are parallel or intersect in a line. If they intersect, you can determine the angle $(0 \le \theta \le \pi/2)$ between them from the angle between their normal vectors, as shown in Figure 11.47. Specifically, if vectors \mathbf{n}_1 and \mathbf{n}_2 are normal to two intersecting planes, the angle θ between the normal vectors is equal to the angle between the two planes and is given by

$$\cos \theta = \frac{|\mathbf{n}_1 \cdot \mathbf{n}_2|}{\|\mathbf{n}_1\| \|\mathbf{n}_2\|}.$$
 Angle between two planes

Consequently, two planes with normal vectors \mathbf{n}_1 and \mathbf{n}_2 are

1. *perpendicular* if $\mathbf{n}_1 \cdot \mathbf{n}_2 = 0$.

2. *parallel* if \mathbf{n}_1 is a scalar multiple of \mathbf{n}_2 .





EXAMPLE 4 Finding the Line of Intersection of Two Planes

Find the angle between the two planes given by

x - 2y + z = 0	Equation of plane 1
2x + 3y - 2z = 0	Equation of plane 2

and find parametric equations of their line of intersection (see Figure 11.48).

Solution Normal vectors for the planes are $\mathbf{n}_1 = \langle 1, -2, 1 \rangle$ and $\mathbf{n}_2 = \langle 2, 3, -2 \rangle$. Consequently, the angle between the two planes is determined as follows.

$$\cos \theta = \frac{|\mathbf{n}_1 \cdot \mathbf{n}_2|}{\|\mathbf{n}_1\| \|\mathbf{n}_2\|}$$
$$= \frac{|-6|}{\sqrt{6}\sqrt{17}}$$
$$= \frac{6}{\sqrt{102}}$$
$$\approx 0.59409$$

Cosine of angle between \mathbf{n}_1 and \mathbf{n}_2

This implies that the angle between the two planes is $\theta \approx 53.55^{\circ}$. You can find the line of intersection of the two planes by simultaneously solving the two linear equations representing the planes. One way to do this is to multiply the first equation by -2 and add the result to the second equation.

$$\begin{array}{c} x - 2y + z = 0 \\ 2x + 3y - 2z = 0 \end{array} \longrightarrow \begin{array}{c} -2x + 4y - 2z = 0 \\ \underline{2x + 3y - 2z = 0} \\ 7y - 4z = 0 \end{array} \longrightarrow \begin{array}{c} y = \frac{4z}{7} \end{array}$$

Substituting y = 4z/7 back into one of the original equations, you can determine that x = z/7. Finally, by letting t = z/7, you obtain the parametric equations

x = t, y = 4t, and z = 7t Line of intersection

which indicate that 1, 4, and 7 are direction numbers for the line of intersection.

Note that the direction numbers in Example 4 can be obtained from the cross product of the two normal vectors as follows.

$$\mathbf{n}_{1} \times \mathbf{n}_{2} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & -2 & 1 \\ 2 & 3 & -2 \end{vmatrix}$$
$$= \begin{vmatrix} -2 & 1 \\ 3 & -2 \end{vmatrix} \mathbf{i} - \begin{vmatrix} 1 & 1 \\ 2 & -2 \end{vmatrix} \mathbf{j} + \begin{vmatrix} 1 & -2 \\ 2 & 3 \end{vmatrix} \mathbf{k}$$
$$= \mathbf{i} + 4\mathbf{i} + 7\mathbf{k}$$

.

This means that the line of intersection of the two planes is parallel to the cross product of their normal vectors.

NOTE The three-dimensional rotatable graphs that are available in the premium eBook for this text can help you visualize surfaces such as those shown in Figure 11.48. If you have access to these graphs, you should use them to help your spatial intuition when studying this section and other sections in the text that deal with vectors, curves, or surfaces in space.

Sketching Planes in Space

If a plane in space intersects one of the coordinate planes, the line of intersection is called the **trace** of the given plane in the coordinate plane. To sketch a plane in space, it is helpful to find its points of intersection with the coordinate axes and its traces in the coordinate planes. For example, consider the plane given by

3x + 2y + 4z = 12. Equation of plane

You can find the *xy*-trace by letting z = 0 and sketching the line

3x + 2y = 12 xy-trace

in the *xy*-plane. This line intersects the *x*-axis at (4, 0, 0) and the *y*-axis at (0, 6, 0). In Figure 11.49, this process is continued by finding the *yz*-trace and the *xz*-trace, and then shading the triangular region lying in the first octant.



z Plane: 2x + z = 1(0, 0, 1) $(\frac{1}{2}, 0, 0)$ y

Plane 2x + z = 1 is parallel to the y-axis. Figure 11.50

If an equation of a plane has a missing variable, such as 2x + z = 1, the plane must be *parallel to the axis* represented by the missing variable, as shown in Figure 11.50. If two variables are missing from an equation of a plane, it is *parallel to the coordinate plane* represented by the missing variables, as shown in Figure 11.51.







Plane ax + d = 0 is parallel to the *yz*-plane **Figure 11.51**

Plane by + d = 0 is parallel to the *xz*-plane

Plane cz + d = 0 is parallel to the *xy*-plane



The distance between a point and a plane **Figure 11.52**

Distances Between Points, Planes, and Lines

This section is concluded with the following discussion of two basic types of problems involving distance in space.

- 1. Finding the distance between a point and a plane
- 2. Finding the distance between a point and a line

The solutions of these problems illustrate the versatility and usefulness of vectors in coordinate geometry: the first problem uses the *dot product* of two vectors, and the second problem uses the *cross product*.

The distance *D* between a point *Q* and a plane is the length of the shortest line segment connecting *Q* to the plane, as shown in Figure 11.52. If *P* is *any* point in the plane, you can find this distance by projecting the vector \overrightarrow{PQ} onto the normal vector **n**. The length of this projection is the desired distance.

THEOREM 11.13 DISTANCE BETWEEN A POINT AND A PLANE

The distance between a plane and a point Q (not in the plane) is

$$D = \|\operatorname{proj}_{\mathbf{n}} \overrightarrow{PQ}\| = \frac{|\overrightarrow{PQ} \cdot \mathbf{n}|}{\|\mathbf{n}\|}$$

where P is a point in the plane and **n** is normal to the plane.

To find a point in the plane given by ax + by + cz + d = 0 ($a \neq 0$), let y = 0 and z = 0. Then, from the equation ax + d = 0, you can conclude that the point (-d/a, 0, 0) lies in the plane.

EXAMPLE 5 Finding the Distance Between a Point and a Plane

Find the distance between the point Q(1, 5, -4) and the plane given by

3x - y + 2z = 6.

Solution You know that $\mathbf{n} = \langle 3, -1, 2 \rangle$ is normal to the given plane. To find a point in the plane, let y = 0 and z = 0, and obtain the point P(2, 0, 0). The vector from P to Q is given by

$$\overline{PQ} = \langle 1 - 2, 5 - 0, -4 - 0 \rangle = \langle -1, 5, -4 \rangle.$$

Using the Distance Formula given in Theorem 11.13 produces

$$D = \frac{|\overline{PQ} \cdot \mathbf{n}|}{\|\mathbf{n}\|} = \frac{|\langle -1, 5, -4 \rangle \cdot \langle 3, -1, 2 \rangle|}{\sqrt{9 + 1 + 4}}$$
Distance between a point and a plane
$$= \frac{|-3 - 5 - 8|}{\sqrt{14}}$$
$$= \frac{16}{\sqrt{14}}.$$

NOTE The choice of the point *P* in Example 5 is arbitrary. Try choosing a different point in the plane to verify that you obtain the same distance.

Ochap

From Theorem 11.13, you can determine that the distance between the point $Q(x_0, y_0, z_0)$ and the plane given by ax + by + cz + d = 0 is

$$D = \frac{|a(x_0 - x_1) + b(y_0 - y_1) + c(z_0 - z_1)|}{\sqrt{a^2 + b^2 + c^2}}$$

or

$$D = \frac{|ax_0 + by_0 + cz_0 + d|}{\sqrt{a^2 + b^2 + c^2}}$$
 Distance between a point and a plane

where $P(x_1, y_1, z_1)$ is a point in the plane and $d = -(ax_1 + by_1 + cz_1)$.

EXAMPLE 6 Finding the Distance Between Two Parallel Planes

Find the distance between the two parallel planes given by

3x - y + 2z - 6 = 0 and 6x - 2y + 4z + 4 = 0.

Solution The two planes are shown in Figure 11.53. To find the distance between the planes, choose a point in the first plane, say $(x_0, y_0, z_0) = (2, 0, 0)$. Then, from the second plane, you can determine that a = 6, b = -2, c = 4, and d = 4, and conclude that the distance is

$$D = \frac{|ax_0 + by_0 + cz_0 + d|}{\sqrt{a^2 + b^2 + c^2}}$$

Distance between a point and a plane
$$= \frac{|6(2) + (-2)(0) + (4)(0) + 4|}{\sqrt{6^2 + (-2)^2 + 4^2}}$$

$$= \frac{16}{\sqrt{56}} = \frac{8}{\sqrt{14}} \approx 2.14.$$

The formula for the distance between a point and a line in space resembles that for the distance between a point and a plane—except that you replace the dot product with the length of the cross product and the normal vector \mathbf{n} with a direction vector for the line.

THEOREM 11.14 DISTANCE BETWEEN A POINT AND A LINE IN SPACE

The distance between a point Q and a line in space is given by

$$D = \frac{\|\overline{PQ} \times \mathbf{u}\|}{\|\mathbf{u}\|}$$

where \mathbf{u} is a direction vector for the line and P is a point on the line.



The distance between a point and a line **Figure 11.54**

PROOF In Figure 11.54, let *D* be the distance between the point *Q* and the given line. Then
$$D = \|\overrightarrow{PQ}\| \sin \theta$$
, where θ is the angle between **u** and \overrightarrow{PQ} . By Property 2 of Theorem 11.8, you have

$$\|\mathbf{u}\| \|\overline{PQ}\| \sin \theta = \|\mathbf{u} \times \overline{PQ}\| = \|\overline{PQ} \times \mathbf{u}\|.$$

Consequently,

$$D = \|\overrightarrow{PQ}\| \sin \theta = \frac{\|\overrightarrow{PQ} \times \mathbf{u}\|}{\|\mathbf{u}\|}.$$

Copyright 2010 Cengage Learning, Inc. All Rights Reserved. May not be copied, scanned, or duplicated, in whole or in part.



The distance between the parallel planes is

approximately 2.14. Figure 11.53

EXAMPLE 7 Finding the Distance Between a Point and a Line

Find the distance between the point Q(3, -1, 4) and the line given by

$$x = -2 + 3t$$
, $y = -2t$, and $z = 1 + 4t$.

Solution Using the direction numbers 3, -2, and 4, you know that a direction vector for the line is

$$\mathbf{u} = \langle 3, -2, 4 \rangle$$
. Direction vector for line

To find a point on the line, let t = 0 and obtain

$$P = (-2, 0, 1).$$
 Point on the line

So,

$$\overrightarrow{PQ} = \langle 3 - (-2), -1 - 0, 4 - 1 \rangle = \langle 5, -1, 3 \rangle$$

.

and you can form the cross product

.

$$\overrightarrow{PQ} \times \mathbf{u} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 5 & -1 & 3 \\ 3 & -2 & 4 \end{vmatrix} = 2\mathbf{i} - 11\mathbf{j} - 7\mathbf{k} = \langle 2, -11, -7 \rangle.$$

Finally, using Theorem 11.14, you can find the distance to be

$$D = \frac{\|\overline{PQ} \times \mathbf{u}\|}{\|\mathbf{u}\|}$$
$$= \frac{\sqrt{174}}{\sqrt{29}}$$
$$= \sqrt{6} \approx 2.45.$$
 See Figure 11.55.



The distance between the point Q and the line is $\sqrt{6} \approx 2.45$. Figure 11.55

11.5 Exercises

See www.CalcChat.com for worked-out solutions to odd-numbered exercises.

In Exercises 1 and 2, the figure shows the graph of a line given by the parametric equations. (a) Draw an arrow on the line to indicate its orientation. To print an enlarged copy of the graph, go to the website *www.mathgraphs.com*. (b) Find the coordinates of two points, *P* and *Q*, on the line. Determine the vector \overrightarrow{PQ} . What is the relationship between the components of the vector and the coefficients of *t* in the parametric equations? Why is this true? (c) Determine the coordinates of any points of intersection with the coordinate planes. If the line does not intersect a coordinate plane, explain why.



In Exercises 3 and 4, determine whether each point lies on the line.

3.
$$x = -2 + t$$
, $y = 3t$, $z = 4 + t$
(a) (0, 6, 6) (b) (2, 3, 5)
4. $\frac{x-3}{2} = \frac{y-7}{8} = z + 2$
(a) (7, 23, 0) (b) (1, -1, -3)

In Exercises 5–10, find sets of (a) parametric equations and (b) symmetric equations of the line through the point parallel to the given vector or line (if possible). (For each line, write the direction numbers as integers.)

Point	Parallel to
5. (0, 0, 0)	$\mathbf{v} = \langle 3, 1, 5 \rangle$
6. (0, 0, 0)	$\mathbf{v} = \left\langle -2, \frac{5}{2}, 1 \right\rangle$
7. (−2, 0, 3)	$\mathbf{v} = 2\mathbf{i} + 4\mathbf{j} - 2\mathbf{k}$
8. (−3, 0, 2)	$\mathbf{v} = 6\mathbf{j} + 3\mathbf{k}$
9. (1, 0, 1)	x = 3 + 3t, y = 5 - 2t, z = -7 + t
10. (-3, 5, 4)	$\frac{x-1}{3} = \frac{y+1}{-2} = z - 3$

In Exercises 11–14, find sets of (a) parametric equations and (b) symmetric equations of the line through the two points (if possible). (For each line, write the direction numbers as integers.)

11. $(5, -3, -2), \left(-\frac{2}{3}, \frac{2}{3}, 1\right)$	12. (0, 4, 3), (-1, 2, 5)
13. $(7, -2, 6), (-3, 0, 6)$	14. (0, 0, 25), (10, 10, 0)

In Exercises 15–22, find a set of parametric equations of the line.

- **15.** The line passes through the point (2, 3, 4) and is parallel to the *xz*-plane and the *yz*-plane.
- 16. The line passes through the point (-4, 5, 2) and is parallel to the *xy*-plane and the *yz*-plane.
- 17. The line passes through the point (2, 3, 4) and is perpendicular to the plane given by 3x + 2y z = 6.
- 18. The line passes through the point (-4, 5, 2) and is perpendicular to the plane given by -x + 2y + z = 5.
- **19.** The line passes through the point (5, -3, -4) and is parallel to $\mathbf{v} = \langle 2, -1, 3 \rangle$.
- **20.** The line passes through the point (-1, 4, -3) and is parallel to $\mathbf{v} = 5\mathbf{i} \mathbf{j}$.
- **21.** The line passes through the point (2, 1, 2) and is parallel to the line x = -t, y = 1 + t, z = -2 + t.
- 22. The line passes through the point (-6, 0, 8) and is parallel to the line x = 5 2t, y = -4 + 2t, z = 0.

In Exercises 23–26, find the coordinates of a point P on the line and a vector v parallel to the line.

23. x = 3 - t, y = -1 + 2t, z = -2 **24.** x = 4t, y = 5 - t, z = 4 + 3t**25.** $\frac{x - 7}{4} = \frac{y + 6}{2} = z + 2$ **26.** $\frac{x + 3}{5} = \frac{y}{8} = \frac{z - 3}{6}$

In Exercises 27–30, determine if any of the lines are parallel or identical.

27.
$$L_1: x = 6 - 3t, y = -2 + 2t, z = 5 + 4t$$

 $L_2: x = 6t, y = 2 - 4t, z = 13 - 8t$
 $L_3: x = 10 - 6t, y = 3 + 4t, z = 7 + 8t$
 $L_4: x = -4 + 6t, y = 3 + 4t, z = 5 - 6t$
28. $L_1: x = 3 + 2t, y = -6t, z = 1 - 2t$
 $L_2: x = 1 + 2t, y = -1 - t, z = 3t$
 $L_3: x = -1 + 2t, y = 3 - 10t, z = 1 - 4t$
 $L_4: x = 5 + 2t, y = 1 - t, z = 8 + 3t$
29. $L_1: \frac{x - 8}{4} = \frac{y + 5}{-2} = \frac{z + 9}{3}$
 $L_2: \frac{x + 7}{2} = \frac{y - 4}{1} = \frac{z + 6}{5}$
 $L_3: \frac{x + 4}{-8} = \frac{y - 1}{4} = \frac{z + 18}{-6}$
 $L_4: \frac{x - 2}{-2} = \frac{y + 3}{1} = \frac{z - 4}{15}$

30.
$$L_1: \frac{x-3}{2} = \frac{y-2}{1} = \frac{z+2}{2}$$

 $L_2: \frac{x-1}{4} = \frac{y-1}{2} = \frac{z+3}{4}$
 $L_3: \frac{x+2}{1} = \frac{y-1}{0.5} = \frac{z-3}{1}$
 $L_4: \frac{x-3}{2} = \frac{y+1}{4} = \frac{z-2}{-1}$

-

- - -

In Exercises 31–34, determine whether the lines intersect, and if so, find the point of intersection and the cosine of the angle of intersection.

31.
$$x = 4t + 2$$
, $y = 3$, $z = -t + 1$
 $x = 2s + 2$, $y = 2s + 3$, $z = s + 1$
32. $x = -3t + 1$, $y = 4t + 1$, $z = 2t + 4$
 $x = 3s + 1$, $y = 2s + 4$, $z = -s + 1$
33. $\frac{x}{3} = \frac{y - 2}{-1} = z + 1$, $\frac{x - 1}{4} = y + 2 = \frac{z + 3}{-3}$
34. $\frac{x - 2}{-3} = \frac{y - 2}{6} = z - 3$, $\frac{x - 3}{2} = y + 5 = \frac{z + 2}{4}$

CAS In Exercises 35 and 36, use a computer algebra system to graph the pair of intersecting lines and find the point of intersection.

35.
$$x = 2t + 3, y = 5t - 2, z = -t + 1$$

 $x = -2s + 7, y = s + 8, z = 2s - 1$
36. $x = 2t - 1, y = -4t + 10, z = t$
 $x = -5s - 12, y = 3s + 11, z = -2s - 4$

Cross Product In Exercises 37 and 38, (a) find the coordinates of three points P, Q, and R in the plane, and determine the vectors \overrightarrow{PQ} and \overrightarrow{PR} . (b) Find $\overrightarrow{PQ} \times \overrightarrow{PR}$. What is the relationship between the components of the cross product and the coefficients of the equation of the plane? Why is this true?



In Exercises 39 and 40, determine whether the plane passes through each point.

39.
$$x + 2y - 4z - 1 = 0$$

(a) $(-7, 2, -1)$ (b) $(5, 2, 2)$
40. $2x + y + 3z - 6 = 0$
(a) $(3, 6, -2)$ (b) $(-1, 5, -1)$

In Exercises 41–46, find an equation of the plane passing through the point perpendicular to the given vector or line.

Point	Perpendicular to
41. (1, 3, -7)	$\mathbf{n} = \mathbf{j}$
42. (0, -1, 4)	$\mathbf{n} = \mathbf{k}$
43. (3, 2, 2)	$\mathbf{n} = 2\mathbf{i} + 3\mathbf{j} - \mathbf{k}$
44. (0, 0, 0)	$\mathbf{n} = -3\mathbf{i} + 2\mathbf{k}$
45. (-1, 4, 0)	x = -1 + 2t, y = 5 - t, z = 3 - 2t
46. (3, 2, 2)	$\frac{x-1}{4} = y + 2 = \frac{z+3}{2}$

In Exercises 47–58, find an equation of the plane.

- **47.** The plane passes through (0, 0, 0), (2, 0, 3), and (-3, -1, 5).
- **48.** The plane passes through (3, -1, 2), (2, 1, 5), and (1, -2, -2).
- **49.** The plane passes through (1, 2, 3), (3, 2, 1), and (-1, -2, 2).
- **50.** The plane passes through the point (1, 2, 3) and is parallel to the *yz*-plane.
- **51.** The plane passes through the point (1, 2, 3) and is parallel to the *xy*-plane.
- **52.** The plane contains the *y*-axis and makes an angle of $\pi/6$ with the positive *x*-axis.
- **53.** The plane contains the lines given by

$$\frac{x-1}{-2} = y - 4 = z$$
 and $\frac{x-2}{-3} = \frac{y-1}{4} = \frac{z-2}{-1}$

54. The plane passes through the point (2, 2, 1) and contains the line given by

 $\frac{x}{2} = \frac{y-4}{-1} = z.$

- **55.** The plane passes through the points (2, 2, 1) and (-1, 1, -1) and is perpendicular to the plane 2x 3y + z = 3.
- 56. The plane passes through the points (3, 2, 1) and (3, 1, -5) and is perpendicular to the plane 6x + 7y + 2z = 10.
- 57. The plane passes through the points (1, -2, -1) and (2, 5, 6) and is parallel to the *x*-axis.
- **58.** The plane passes through the points (4, 2, 1) and (-3, 5, 7) and is parallel to the *z*-axis.

In Exercises 59 and 60, sketch a graph of the line and find the points (if any) where the line intersects the *xy*-, *xz*-, and *yz*-planes.

t

59.
$$x = 1 - 2t$$
, $y = -2 + 3t$, $z = -4 + 60$. $\frac{x - 2}{3} = y + 1 = \frac{z - 3}{2}$

In Exercises 61–64, find an equation of the plane that contains all the points that are equidistant from the given points.

61. (2, 2, 0), (0, 2, 2)**62.** (1, 0, 2), (2, 0, 1)**63.** (-3, 1, 2), (6, -2, 4)**64.** (-5, 1, -3), (2, -1, 6)

In Exercises 65–70, determine whether the planes are parallel, orthogonal, or neither. If they are neither parallel nor orthogonal, find the angle of intersection.

65.	5x - 3y + z = 4	66.	3x + y - 4z = 3
	x + 4y + 7z = 1		-9x - 3y + 12z = 4
67.	x - 3y + 6z = 4	68.	3x + 2y - z = 7
	5x + y - z = 4		x - 4y + 2z = 0
69.	x - 5y - z = 1	70.	2x - z = 1
	5x - 25y - 5z = -3		4x + y + 8z = 10

In Exercises 71–78, sketch a graph of the plane and label any intercepts.

71 $4x + 2y + 6z = 12$	72. $3x + 6y + 2z = 6$
73. $2x - y + 3z = 4$	74. $2x - y + z = 4$
75. $x + z = 6$	76. $2x + y = 8$
77. $x = 5$	78. $z = 8$

CAS In Exercises 79–82, use a computer algebra system to graph the plane.

79. $2x + y - z = 6$	80. $x - 3z = 3$
81. $-5x + 4y - 6z = -8$	82. $2.1x - 4.7y - z = -3$

In Exercises 83–86, determine if any of the planes are parallel or identical.

83. $P_1: 15x - 6y + 24z = 17$ $P_2: -5x + 2y - 8z = 6$ $P_3: 6x - 4y + 4z = 9$ $P_4: 3x - 2y - 2z = 4$ **85.** $P_1: 3x - 2y + 5z = 10$ $P_2: -6x + 4y - 10z = 5$ $P_3: -3x + 2y + 5z = 8$ $P_4: 75x - 50y + 125z = 250$ **86.** $P_1: -60x + 90y + 30z = 27$ $P_2: 6x - 9y - 3z = 2$ $P_3: -20x + 30y + 10z = 9$ $P_4: 12x - 18y + 6z = 5$

In Exercises 87–90, describe the family of planes represented by the equation, where *c* is any real number.

87.	x + y + z = c	88.	x +	y = c
89.	cy + z = 0	90.	<i>x</i> +	cz = 0

In Exercises 91 and 92, (a) find the angle between the two planes, and (b) find a set of parametric equations for the line of intersection of the planes.

91. $3x + 2y - z = 7$	92. $6x - 3y + z = 5$
x - 4y + 2z = 0	-x + y + 5z = 5

In Exercises 93-96, find the point(s) of intersection (if any) of the plane and the line. Also determine whether the line lies in the plane.

93.
$$2x - 2y + z = 12$$
, $x - \frac{1}{2} = \frac{y + (3/2)}{-1} = \frac{z + 1}{2}$
94. $2x + 3y = -5$, $\frac{x - 1}{4} = \frac{y}{2} = \frac{z - 3}{6}$
95. $2x + 3y = 10$, $\frac{x - 1}{3} = \frac{y + 1}{-2} = z - 3$
96. $5x + 3y = 17$, $\frac{x - 4}{2} = \frac{y + 1}{-3} = \frac{z + 2}{5}$

In Exercises 97–100, find the distance between the point and the plane.

97.
$$(0, 0, 0)$$
98. $(0, 0, 0)$ $2x + 3y + z = 12$ $5x + y - z = 9$ **99.** $(2, 8, 4)$ **100.** $(1, 3, -1)$ $2x + y + z = 5$ $3x - 4y + 5z = 6$

In Exercises 101–104, verify that the two planes are parallel, and find the distance between the planes.

101. $x - 3y + 4z = 10$	102. $4x - 4y + 9z = 7$
x - 3y + 4z = 6	4x - 4y + 9z = 18
103. $-3x + 6y + 7z = 1$	104. $2x - 4z = 4$
6x - 12y - 14z = 25	2x - 4z = 10

In Exercises 105–108, find the distance between the point and the line given by the set of parametric equations.

105. $(1, 5, -2);$	x = 4t - 2, y = 3,	z = -t + 1
106. $(1, -2, 4);$	x = 2t, y = t - 3,	z = 2t + 2
107. (-2, 1, 3);	x = 1 - t, y = 2 +	t, z = -2t
108. (4, -1, 5);	x = 3, y = 1 + 3t,	z = 1 + t

In Exercises 109 and 110, verify that the lines are parallel, and find the distance between them.

109. $L_1: x = 2 - t$, y = 3 + 2t, z = 4 + t $L_2: x = 3t$, y = 1 - 6t, z = 4 - 3t**110.** $L_1: x = 3 + 6t$, y = -2 + 9t, z = 1 - 12t $L_2: x = -1 + 4t$, y = 3 + 6t, z = -8t

WRITING ABOUT CONCEPTS

- **111.** Give the parametric equations and the symmetric equations of a line in space. Describe what is required to find these equations.
- **112.** Give the standard equation of a plane in space. Describe what is required to find this equation.
- **113.** Describe a method of finding the line of intersection of two planes.
- **114.** Describe each surface given by the equations x = a, y = b, and z = c.

WRITING ABOUT CONCEPTS (continued)

115. Describe a method for determining when two planes

$$a_1x + b_1y + c_1z + d_1 = 0$$
 and
 $a_2x + b_2y + c_2z + d_2 = 0$

are (a) parallel and (b) perpendicular. Explain your reasoning.

- **116.** Let L_1 and L_2 be nonparallel lines that do not intersect. Is it possible to find a nonzero vector **v** such that **v** is perpendicular to both L_1 and L_2 ? Explain your reasoning.
- **117.** Find an equation of the plane with *x*-intercept (*a*, 0, 0), *y*-intercept (0, *b*, 0), and *z*-intercept (0, 0, *c*). (Assume *a*, *b*, and *c* are nonzero.)

CAPSTONE

- **118.** Match the equation or set of equations with the description it represents.
 - (a) Set of parametric equations of a line
 - (b) Set of symmetric equations of a line
 - (c) Standard equation of a plane in space
 - (d) General form of an equation of a plane in space
 - (i) (x 6)/2 = (y + 1)/-3 = z/1
 - (ii) 2x 7y + 5z + 10 = 0
 - (iii) x = 4 + 7t, y = 3 + t, z = 3 3t
 - (iv) 2(x 1) + (y + 3) 4(z 5) = 0
- **119.** Describe and find an equation for the surface generated by all points (x, y, z) that are four units from the point (3, -2, 5).
- 120. Describe and find an equation for the surface generated by all points (x, y, z) that are four units from the plane 4x 3y + z = 10.
- **121.** *Modeling Data* Per capita consumptions (in gallons) of different types of milk in the United States from 1999 through 2005 are shown in the table. Consumptions of flavored milk, plain reduced-fat milk, and plain light and skim milks are represented by the variables *x*, *y*, and *z*, respectively. *(Source: U.S. Department of Agriculture)*

Year	1999	2000	2001	2002	2003	2004	2005
x	1.4	1.4	1.4	1.6	1.6	1.7	1.7
у	7.3	7.1	7.0	7.0	6.9	6.9	6.9
z	6.2	6.1	5.9	5.8	5.6	5.5	5.6

A model for the data is given by 0.92x - 1.03y + z = 0.02.

- (a) Complete a fourth row in the table using the model to approximate *z* for the given values of *x* and *y*. Compare the approximations with the actual values of *z*.
- (b) According to this model, any increases in consumption of two types of milk will have what effect on the consumption of the third type?

122. *Mechanical Design* The figure shows a chute at the top of a grain elevator of a combine that funnels the grain into a bin. Find the angle between two adjacent sides.



123. *Distance* Two insects are crawling along different lines in three-space. At time t (in minutes), the first insect is at the point (x, y, z) on the line x = 6 + t, y = 8 - t, z = 3 + t. Also, at time t, the second insect is at the point (x, y, z) on the line x = 1 + t, y = 2 + t, z = 2t.

Assume that distances are given in inches.

- (a) Find the distance between the two insects at time t = 0.
- (b) Use a graphing utility to graph the distance between the insects from t = 0 to t = 10.
 - (c) Using the graph from part (b), what can you conclude about the distance between the insects?
 - (d) How close to each other do the insects get?

SECTION PROJECT

Distances in Space

You have learned two distance formulas in this section—the distance between a point and a plane, and the distance between a point and a line. In this project you will study a third distance problem—the distance between two skew lines. Two lines in space are *skew* if they are neither parallel nor intersecting (see figure).

(a) Consider the following two lines in space.

$$L_1: x = 4 + 5t, y = 5 + 5t, z = 1 - 4t$$

- $L_2: x = 4 + s, y = -6 + 8s, z = 7 3s$
- (i) Show that these lines are not parallel.
- (ii) Show that these lines do not intersect, and therefore are skew lines.
- (iii) Show that the two lines lie in parallel planes.
- (iv) Find the distance between the parallel planes from part(iii). This is the distance between the original skew lines.

+ s

(b) Use the procedure in part (a) to find the distance between the lines.

$$L_1: x = 2t, y = 4t, z = 6t$$
$$L_2: x = 1 - s, y = 4 + s, z = -1$$

- **124.** Find the standard equation of the sphere with center (-3, 2, 4) that is tangent to the plane given by 2x + 4y 3z = 8.
- **125.** Find the point of intersection of the plane 3x y + 4z = 7 and the line through (5, 4, -3) that is perpendicular to this plane.
- **126.** Show that the plane 2x y 3z = 4 is parallel to the line x = -2 + 2t, y = -1 + 4t, z = 4, and find the distance between them.
- **127.** Find the point of intersection of the line through (1, -3, 1) and (3, -4, 2), and the plane given by x y + z = 2.
- **128.** Find a set of parametric equations for the line passing through the point (1, 0, 2) that is parallel to the plane given by x + y + z = 5, and perpendicular to the line x = t, y = 1 + t, z = 1 + t.

True or False? In Exercises 129–134, determine whether the statement is true or false. If it is false, explain why or give an example that shows it is false.

- **129.** If $\mathbf{v} = a_1\mathbf{i} + b_1\mathbf{j} + c_1\mathbf{k}$ is any vector in the plane given by $a_2x + b_2y + c_2z + d_2 = 0$, then $a_1a_2 + b_1b_2 + c_1c_2 = 0$.
- 130. Every two lines in space are either intersecting or parallel.
- 131. Two planes in space are either intersecting or parallel.
- **132.** If two lines L_1 and L_2 are parallel to a plane *P*, then L_1 and L_2 are parallel.
- 133. Two planes perpendicular to a third plane in space are parallel.
- 134. A plane and a line in space are either intersecting or parallel.

(c) Use the procedure in part (a) to find the distance between the lines.

$$L_1: x = 3t, y = 2 - t, z = -1 + t$$

- $L_2: x = 1 + 4s, y = -2 + s, z = -3 3s$
- (d) Develop a formula for finding the distance between the skew lines.

$$L_1: x = x_1 + a_1t, \ y = y_1 + b_1t, \ z = z_1 + c_1t$$
$$L_2: x = x_2 + a_2s, \ y = y_2 + b_2s, \ z = z_2 + c_2s$$



11.6 Surfaces in Space



Rulings are parallel to *z*-axis. **Figure 11.56**



Cylinder: Rulings intersect *C* and are parallel to the given line. **Figure 11.57**

- Recognize and write equations of cylindrical surfaces.
- Recognize and write equations of quadric surfaces.
- Recognize and write equations of surfaces of revolution.

Cylindrical Surfaces

The first five sections of this chapter contained the vector portion of the preliminary work necessary to study vector calculus and the calculus of space. In this and the next section, you will study surfaces in space and alternative coordinate systems for space. You have already studied two special types of surfaces.

1.	Spheres: $(x - x_0)^2 + (y - y_0)^2 + (z - z_0)^2 = r^2$	Section 11.2
2.	Planes: $ax + by + cz + d = 0$	Section 11.5

A third type of surface in space is called a **cylindrical surface**, or simply a **cylinder**. To define a cylinder, consider the familiar right circular cylinder shown in Figure 11.56. You can imagine that this cylinder is generated by a vertical line moving around the circle $x^2 + y^2 = a^2$ in the *xy*-plane. This circle is called a **generating curve** for the cylinder, as indicated in the following definition.

DEFINITION OF A CYLINDER

Let C be a curve in a plane and let L be a line not in a parallel plane. The set of all lines parallel to L and intersecting C is called a **cylinder**. C is called the **generating curve** (or **directrix**) of the cylinder, and the parallel lines are called **rulings**.

NOTE Without loss of generality, you can assume that C lies in one of the three coordinate planes. Moreover, this text restricts the discussion to *right* cylinders—cylinders whose rulings are perpendicular to the coordinate plane containing C, as shown in Figure 11.57.

For the right circular cylinder shown in Figure 11.56, the equation of the generating curve is

$$x^2 + y^2 = a^2$$

Equation of generating curve in xy-plane

To find an equation of the cylinder, note that you can generate any one of the rulings by fixing the values of x and y and then allowing z to take on all real values. In this sense, the value of z is arbitrary and is, therefore, not included in the equation. In other words, the equation of this cylinder is simply the equation of its generating curve.

$$x^2 + y^2 = a^2$$

Equation of cylinder in space

EQUATIONS OF CYLINDERS

The equation of a cylinder whose rulings are parallel to one of the coordinate axes contains only the variables corresponding to the other two axes.



Sketch the surface represented by each equation.

a.
$$z = y^2$$
 b. $z = \sin x$, $0 \le x \le 2\pi$

Solution

- **a.** The graph is a cylinder whose generating curve, $z = y^2$, is a parabola in the yz-plane. The rulings of the cylinder are parallel to the x-axis, as shown in Figure 11.58(a).
- **b.** The graph is a cylinder generated by the sine curve in the *xz*-plane. The rulings are parallel to the y-axis, as shown in Figure 11.58(b).



Figure 11.58

STUDY TIP In the table on pages 814

and 815, only one of several orientations of each quadric surface is shown. If the surface is oriented along a different axis, its standard equation will change accordingly, as illustrated in Examples 2 and 3. The fact that the two types of paraboloids have one variable raised to the first power can be helpful in classifying quadric surfaces. The other four types of basic quadric surfaces have equations that are of second degree in all three variables.

Quadric Surfaces

The fourth basic type of surface in space is a **quadric surface**. Ouadric surfaces are the three-dimensional analogs of conic sections.

QUADRIC SURFACE

The equation of a quadric surface in space is a second-degree equation in three variables. The general form of the equation is

 $Ax^{2} + By^{2} + Cz^{2} + Dxy + Exz + Fyz + Gx + Hy + Iz + J = 0.$

There are six basic types of quadric surfaces: ellipsoid, hyperboloid of one sheet, hyperboloid of two sheets, elliptic cone, elliptic paraboloid, and hyperbolic paraboloid.

The intersection of a surface with a plane is called the trace of the surface in the plane. To visualize a surface in space, it is helpful to determine its traces in some wellchosen planes. The traces of quadric surfaces are conics. These traces, together with the standard form of the equation of each quadric surface, are shown in the table on pages 814 and 815.





To classify a quadric surface, begin by writing the surface in standard form. Then, determine several traces taken in the coordinate planes *or* taken in planes that are parallel to the coordinate planes.

EXAMPLE 2 Sketching a Quadric Surface

Classify and sketch the surface given by $4x^2 - 3y^2 + 12z^2 + 12 = 0$.

Solution Begin by writing the equation in standard form.

$4x^2 - 3y^2 + 12z^2 + 12 = 0$	Write original equation.
$\frac{x^2}{-3} + \frac{y^2}{4} - z^2 - 1 = 0$	Divide by -12 .
$\frac{y^2}{4} - \frac{x^2}{3} - \frac{z^2}{1} = 1$	Standard form

From the table on pages 814 and 815, you can conclude that the surface is a hyperboloid of two sheets with the *y*-axis as its axis. To sketch the graph of this surface, it helps to find the traces in the coordinate planes.

xy-trace $(z = 0)$:	$\frac{y^2}{4} - \frac{x^2}{3} = 1$	Hyperbola
<i>xz</i> -trace $(y = 0)$:	$\frac{x^2}{3} + \frac{z^2}{1} = -1$	No trace
yz-trace ($x = 0$):	$\frac{y^2}{4} - \frac{z^2}{1} = 1$	Hyperbola

The graph is shown in Figure 11.59.

EXAMPLE 3 Sketching a Quadric Surface

Classify and sketch the surface given by $x - y^2 - 4z^2 = 0$.

Solution Because x is raised only to the first power, the surface is a paraboloid. The axis of the paraboloid is the x-axis. In the standard form, the equation is

$$x = y^2 + 4z^2$$
. Standard form

Some convenient traces are as follows.

xy-trace $(z = 0)$:	$x = y^2$	Parabola
<i>xz</i> -trace $(y = 0)$:	$x = 4z^2$	Parabola
parallel to yz -plane ($x = 4$):	$\frac{y^2}{4} + \frac{z^2}{1} = 1$	Ellipse

The surface is an *elliptic* paraboloid, as shown in Figure 11.60.

Some second-degree equations in x, y, and z do not represent any of the basic types of quadric surfaces. Here are two examples.

$x^2 + y^2 + z^2 = 0$	Single point
$x^2 + y^2 = 1$	Right circular cylinder











An ellipsoid centered at (2, -1, 1)Figure 11.61

For a quadric surface not centered at the origin, you can form the standard equation by completing the square, as demonstrated in Example 4.

JEXAMPLE 4 A Quadric Surface Not Centered at the Origin

Classify and sketch the surface given by

 $x^2 + 2y^2 + z^2 - 4x + 4y - 2z + 3 = 0.$

Solution Completing the square for each variable produces the following.

 $(x^{2} - 4x +) + 2(y^{2} + 2y +) + (z^{2} - 2z +) = -3$ $(x^{2} - 4x + 4) + 2(y^{2} + 2y + 1) + (z^{2} - 2z + 1) = -3 + 4 + 2 + 1$ $(x - 2)^{2} + 2(y + 1)^{2} + (z - 1)^{2} = 4$ $\frac{(x - 2)^{2}}{4} + \frac{(y + 1)^{2}}{2} + \frac{(z - 1)^{2}}{4} = 1$

From this equation, you can see that the quadric surface is an ellipsoid that is centered at (2, -1, 1). Its graph is shown in Figure 11.61.

TECHNOLOGY A computer algebra system can help you visualize a surface in space.* Most of these computer algebra systems create three-dimensional illusions by sketching several traces of the surface and then applying a "hidden-line" routine that blocks out portions of the surface that lie behind other portions of the surface. Two examples of figures that were generated by *Mathematica* are shown below.



Using a graphing utility to graph a surface in space requires practice. For one thing, you must know enough about the surface to be able to specify a *viewing window* that gives a representative view of the surface. Also, you can often improve the view of a surface by rotating the axes. For instance, note that the elliptic paraboloid in the figure is seen from a line of sight that is "higher" than the line of sight used to view the hyperbolic paraboloid.

*Some 3-D graphing utilities require surfaces to be entered with parametric equations. For a discussion of this technique, see Section 15.5.



Figure 11.62

Surfaces of Revolution

The fifth special type of surface you will study is called a **surface of revolution.** In Section 7.4, you studied a method for finding the *area* of such a surface. You will now look at a procedure for finding its equation. Consider the graph of the radius function

$$y = r(z)$$

Generating curve

in the *yz*-plane. If this graph is revolved about the *z*-axis, it forms a surface of revolution, as shown in Figure 11.62. The trace of the surface in the plane $z = z_0$ is a circle whose radius is $r(z_0)$ and whose equation is

$$x^2 + y^2 = [r(z_0)]^2.$$

Replacing z_0 with z produces an equation that is valid for all values of z. In a similar manner, you can obtain equations for surfaces of revolution for the other two axes, and the results are summarized as follows.

Circular trace in plane: $z = z_0$

SURFACE OF REVOLUTION

If the graph of a radius function r is revolved about one of the coordinate axes, the equation of the resulting surface of revolution has one of the following forms.

- 1. Revolved about the x-axis: $y^2 + z^2 = [r(x)]^2$
- 2. Revolved about the y-axis: $x^2 + z^2 = [r(y)]^2$
- 3. Revolved about the z-axis: $x^2 + y^2 = [r(z)]^2$

EXAMPLE 5 Finding an Equation for a Surface of Revolution

a. An equation for the surface of revolution formed by revolving the graph of







$$y = \frac{1}{z}$$
Radius function
ut the z-axis is

$$x^{2} + y^{2} = [r(z)]^{2}$$
Revolved about the z-axis

$$x^{2} + y^{2} = \left(\frac{1}{z}\right)^{2}.$$
Substitute 1/z for r(z).

Substitute 1/z for r(z).

b. To find an equation for the surface formed by revolving the graph of $9x^2 = y^3$ about the y-axis, solve for x in terms of y to obtain

$$=\frac{1}{3}y^{3/2} = r(y).$$

 $y = \frac{1}{7}$

about the z-axis is

х

So, the equation for this surface is

 $x^2 + z^2 = [r(y)]^2$ Revolved about the y-axis $x^2 + z^2 = \left(\frac{1}{3}y^{3/2}\right)^2$ Substitute $\frac{1}{3}y^{3/2}$ for r(y). $x^2 + z^2 = \frac{1}{9}y^3$. Equation of surface

The graph is shown in Figure 11.63.

The generating curve for a surface of revolution is not unique. For instance, the surface

$$x^2 + z^2 = e^{-2y}$$

can be formed by revolving either the graph of $x = e^{-y}$ about the y-axis or the graph of $z = e^{-y}$ about the y-axis, as shown in Figure 11.64.



Figure 11.64

EXAMPLE 6 Finding a Generating Curve for a Surface of Revolution

Find a generating curve and the axis of revolution for the surface given by

$$x^2 + 3y^2 + z^2 = 9.$$

Solution You now know that the equation has one of the following forms.

$$x^{2} + y^{2} = [r(z)]^{2}$$
Revolved about *z*-axis
$$y^{2} + z^{2} = [r(x)]^{2}$$
Revolved about *x*-axis
$$x^{2} + z^{2} = [r(y)]^{2}$$
Revolved about *y*-axis

Because the coefficients of x^2 and z^2 are equal, you should choose the third form and write

$$x^2 + z^2 = 9 - 3y^2.$$

The *y*-axis is the axis of revolution. You can choose a generating curve from either of the following traces.

$x^2 = 9 - 3y^2$	Trace in <i>xy</i> -plane
$z^2 = 9 - 3y^2$	Trace in yz-plane

For example, using the first trace, the generating curve is the semiellipse given by

 $x = \sqrt{9 - 3y^2}$. Generating curve

The graph of this surface is shown in Figure 11.65.





11.6 Exercises

See www.CalcChat.com for worked-out solutions to odd-numbered exercises.

In Exercises 1–6, match the equation with its graph. [The graphs are labeled (a), (b), (c), (d), (e), and (f).]



In Exercises 7–16, describe and sketch the surface.

7. $y = 5$	8. $z = 2$
9. $y^2 + z^2 = 9$	10. $x^2 + z^2 = 25$
11. $x^2 - y = 0$	12. $y^2 + z = 6$
13. $4x^2 + y^2 = 4$	14. $y^2 - z^2 = 16$
15. $z - \sin y = 0$	16. $z - e^y = 0$

17. *Think About It* The four figures are graphs of the quadric surface $z = x^2 + y^2$. Match each of the four graphs with the point in space from which the paraboloid is viewed. The four points are (0, 0, 20), (0, 20, 0), (20, 0, 0), and (10, 10, 20).







- **CAS** 18. Use a computer algebra system to graph a view of the cylinder $y^2 + z^2 = 4$ from each point.
 - (a) (10, 0, 0)
 - (b) (0, 10, 0)
 - (c) (10, 10, 10)

In Exercises 19–32, identify and sketch the quadric surface. Use a computer algebra system to confirm your sketch.

19. $x^2 + \frac{y^2}{4} + z^2 = 1$	20. $\frac{x^2}{16} + \frac{y^2}{25} + \frac{z^2}{25} = 1$
21. $16x^2 - y^2 + 16z^2 = 4$	22. $-8x^2 + 18y^2 + 18z^2 = 2$
23. $4x^2 - y^2 - z^2 = 1$	24. $z^2 - x^2 - \frac{y^2}{4} = 1$
25. $x^2 - y + z^2 = 0$	26. $z = x^2 + 4y^2$
27. $x^2 - y^2 + z = 0$	28. $3z = -y^2 + x^2$
29. $z^2 = x^2 + \frac{y^2}{9}$	30. $x^2 = 2y^2 + 2z^2$
31. $16x^2 + 9y^2 + 16z^2 - 32x - $	36y + 36 = 0
32. $9x^2 + y^2 - 9z^2 - 54x - 4y$	-54z + 4 = 0

CAS In Exercises 33–42, use a computer algebra system to graph the surface. (*Hint:* It may be necessary to solve for z and acquire two equations to graph the surface.)

33. $z = 2 \cos x$	34. $z = x^2 + 0.5y^2$
35. $z^2 = x^2 + 7.5y^2$	36. $3.25y = x^2 + z^2$
37. $x^2 + y^2 = \left(\frac{2}{z}\right)^2$	38. $x^2 + y^2 = e^{-z}$
39. $z = 10 - \sqrt{ xy }$	40. $z = \frac{-x}{8 + x^2 + y^2}$
41. $6x^2 - 4y^2 + 6z^2 = -36$	42. $9x^2 + 4y^2 - 8z^2 = 72$

In Exercises 43–46, sketch the region bounded by the graphs of the equations.

43.
$$z = 2\sqrt{x^2 + y^2}$$
, $z = 2$
44. $z = \sqrt{4 - x^2}$, $y = \sqrt{4 - x^2}$, $x = 0$, $y = 0$, $z = 0$
45. $x^2 + y^2 = 1$, $x + z = 2$, $z = 0$
46. $z = \sqrt{4 - x^2 - y^2}$, $y = 2z$, $z = 0$

In Exercises 47–52, find an equation for the surface of revolution generated by revolving the curve in the indicated coordinate plane about the given axis.

	Equation of Curve	Coordinate Plane	Axis of Revolution
47.	$z^2 = 4y$	yz-plane	y-axis
48.	z = 3y	yz-plane	y-axis
49.	z = 2y	yz-plane	z-axis
50.	$2z = \sqrt{4 - x^2}$	xz-plane	<i>x</i> -axis
51.	xy = 2	xy-plane	<i>x</i> -axis
52.	$z = \ln y$	yz-plane	z-axis

In Exercises 53 and 54, find an equation of a generating curve given the equation of its surface of revolution.

53.
$$x^2 + y^2 - 2z = 0$$
 54. $x^2 + z^2 = \cos^2 y$

WRITING ABOUT CONCEPTS

- 55. State the definition of a cylinder.
- **56.** What is meant by the trace of a surface? How do you find a trace?
- **57.** Identify the six quadric surfaces and give the standard form of each.

CAPSTONE

58. What does the equation $z = x^2$ represent in the *xz*-plane? What does it represent in three-space?

In Exercises 59 and 60, use the shell method to find the volume of the solid below the surface of revolution and above the *xy*-plane.

- **59.** The curve $z = 4x x^2$ in the *xz*-plane is revolved about the *z*-axis.
- **60.** The curve $z = \sin y$ ($0 \le y \le \pi$) in the *yz*-plane is revolved about the *z*-axis.

In Exercises 61 and 62, analyze the trace when the surface

 $z = \frac{1}{2}x^2 + \frac{1}{4}y^2$

is intersected by the indicated planes.

61. Find the lengths of the major and minor axes and the coordinates of the foci of the ellipse generated when the surface is intersected by the planes given by

(a) z = 2 and (b) z = 8.

62. Find the coordinates of the focus of the parabola formed when the surface is intersected by the planes given by

(a) y = 4 and (b) x = 2.

In Exercises 63 and 64, find an equation of the surface satisfying the conditions, and identify the surface.

63. The set of all points equidistant from the point (0, 2, 0) and the plane y = -2

- **64.** The set of all points equidistant from the point (0, 0, 4) and the *xy*-plane
- **65.** *Geography* Because of the forces caused by its rotation, Earth is an oblate ellipsoid rather than a sphere. The equatorial radius is 3963 miles and the polar radius is 3950 miles. Find an equation of the ellipsoid. (Assume that the center of Earth is at the origin and that the trace formed by the plane z = 0corresponds to the equator.)
- **66.** *Machine Design* The top of a rubber bushing designed to absorb vibrations in an automobile is the surface of revolution generated by revolving the curve $z = \frac{1}{2}y^2 + 1$ ($0 \le y \le 2$) in the *yz*-plane about the *z*-axis.
 - (a) Find an equation for the surface of revolution.
 - (b) All measurements are in centimeters and the bushing is set on the *xy*-plane. Use the shell method to find its volume.
 - (c) The bushing has a hole of diameter 1 centimeter through its center and parallel to the axis of revolution. Find the volume of the rubber bushing.
- **67.** Determine the intersection of the hyperbolic paraboloid $z = y^2/b^2 x^2/a^2$ with the plane bx + ay z = 0. (Assume a, b > 0.)
- **68.** Explain why the curve of intersection of the surfaces $x^2 + 3y^2 2z^2 + 2y = 4$ and $2x^2 + 6y^2 4z^2 3x = 2$ lies in a plane.

True or False? In Exercises 69–72, determine whether the statement is true or false. If it is false, explain why or give an example that shows it is false.

- **69.** A sphere is an ellipsoid.
- 70. The generating curve for a surface of revolution is unique.
- 71. All traces of an ellipsoid are ellipses.
- 72. All traces of a hyperboloid of one sheet are hyperboloids.
- **73.** *Think About It* Three types of classic "topological" surfaces are shown below. The sphere and torus have both an "inside" and an "outside." Does the Klein bottle have both an inside and an outside? Explain.



11.7 Cylindrical and Spherical Coordinates

- Use cylindrical coordinates to represent surfaces in space.
- Use spherical coordinates to represent surfaces in space.

Cylindrical Coordinates

You have already seen that some two-dimensional graphs are easier to represent in polar coordinates than in rectangular coordinates. A similar situation exists for surfaces in space. In this section, you will study two alternative space-coordinate systems. The first, the **cylindrical coordinate system**, is an extension of polar coordinates in the plane to three-dimensional space.



In a **cylindrical coordinate system,** a point *P* in space is represented by an ordered triple (r, θ, z) .

1. (r, θ) is a polar representation of the projection of *P* in the *xy*-plane.

2. *z* is the directed distance from (r, θ) to *P*.

To convert from rectangular to cylindrical coordinates (or vice versa), use the following conversion guidelines for polar coordinates, as illustrated in Figure 11.66.

Cylindrical to rectangular:

$$x = r \cos \theta$$
, $y = r \sin \theta$, $z = z$

Rectangular to cylindrical:

 $r^2 = x^2 + y^2$, $\tan \theta = \frac{y}{r}$,

The point (0, 0, 0) is called the **pole.** Moreover, because the representation of a point in the polar coordinate system is not unique, it follows that the representation in the cylindrical coordinate system is also not unique.

z = z

EXAMPLE 1 Converting from Cylindrical to Rectangular Coordinates

Convert the point $(r, \theta, z) = \left(4, \frac{5\pi}{6}, 3\right)$ to rectangular coordinates.

Solution Using the cylindrical-to-rectangular conversion equations produces

$$x = 4\cos\frac{5\pi}{6} = 4\left(-\frac{\sqrt{3}}{2}\right) = -2\sqrt{3}$$
$$y = 4\sin\frac{5\pi}{6} = 4\left(\frac{1}{2}\right) = 2$$
$$z = 3.$$

So, in rectangular coordinates, the point is $(x, y, z) = (-2\sqrt{3}, 2, 3)$, as shown in Figure 11.67.



Figure 11.66











EXAMPLE 2 Converting from Rectangular to Cylindrical Coordinates

Convert the point $(x, y, z) = (1, \sqrt{3}, 2)$ to cylindrical coordinates. **Solution** Use the rectangular-to-cylindrical conversion equations.

$$r = \pm \sqrt{1+3} = \pm 2$$

$$\tan \theta = \sqrt{3} \qquad \implies \qquad \theta = \arctan\left(\sqrt{3}\right) + n\pi = \frac{\pi}{3} + n\pi$$

$$z = 2$$

You have two choices for r and infinitely many choices for θ . As shown in Figure 11.68, two convenient representations of the point are

$$\left(2, \frac{\pi}{3}, 2\right) \qquad r > 0 \text{ and } \theta \text{ in Quadrant I}$$

$$\left(-2, \frac{4\pi}{3}, 2\right) \qquad r < 0 \text{ and } \theta \text{ in Quadrant III}$$

Cylindrical coordinates are especially convenient for representing cylindrical surfaces and surfaces of revolution with the *z*-axis as the axis of symmetry, as shown in Figure 11.69.



Vertical planes containing the *z*-axis and horizontal planes also have simple cylindrical coordinate equations, as shown in Figure 11.70.



Figure 11.70











EXAMPLE 3 Rectangular-to-Cylindrical Conversion

Find an equation in cylindrical coordinates for the surface represented by each rectangular equation.

a.
$$x^2 + y^2 = 4z^2$$

b. $y^2 = x$

Solution

a. From the preceding section, you know that the graph $x^2 + y^2 = 4z^2$ is an elliptic cone with its axis along the *z*-axis, as shown in Figure 11.71. If you replace $x^2 + y^2$ with r^2 , the equation in cylindrical coordinates is

$x^2 + y^2 = 4z^2$	Rectangular equation
$r^2 = 4z^2.$	Cylindrical equation

b. The graph of the surface $y^2 = x$ is a parabolic cylinder with rulings parallel to the *z*-axis, as shown in Figure 11.72. By replacing y^2 with $r^2 \sin^2 \theta$ and *x* with $r \cos \theta$, you obtain the following equation in cylindrical coordinates.

$y^2 = x$	Rectangular equation
$r^2\sin^2\theta = r\cos\theta$	Substitute $r \sin \theta$ for y and $r \cos \theta$ for x .
$r(r\sin^2\theta-\cos\theta)=0$	Collect terms and factor.
$r\sin^2\theta - \cos\theta = 0$	Divide each side by <i>r</i> .
$r = \frac{\cos \theta}{\sin^2 \theta}$	Solve for <i>r</i> .
$r = \csc \theta \cot \theta$	Cylindrical equation

Note that this equation includes a point for which r = 0, so nothing was lost by dividing each side by the factor r.

Converting from cylindrical coordinates to rectangular coordinates is less straightforward than converting from rectangular coordinates to cylindrical coordinates, as demonstrated in Example 4.

EXAMPLE 4 Cylindrical-to-Rectangular Conversion

Find an equation in rectangular coordinates for the surface represented by the cylindrical equation

$$r^2\cos 2\theta + z^2 + 1 = 0$$

Solution

 $r^{2} \cos 2\theta + z^{2} + 1 = 0$ $r^{2}(\cos^{2} \theta - \sin^{2} \theta) + z^{2} + 1 = 0$ $r^{2} \cos^{2} \theta - r^{2} \sin^{2} \theta + z^{2} = -1$ $x^{2} - y^{2} + z^{2} = -1$ $y^{2} - x^{2} - z^{2} = 1$ Replace $r \cos \theta$ with x and $r \sin \theta$ with y. Rectangular equation

This is a hyperboloid of two sheets whose axis lies along the *y*-axis, as shown in Figure 11.73.



Figure 11.74

Spherical Coordinates

In the **spherical coordinate system**, each point is represented by an ordered triple: the first coordinate is a distance, and the second and third coordinates are angles. This system is similar to the latitude-longitude system used to identify points on the surface of Earth. For example, the point on the surface of Earth whose latitude is 40° North (of the equator) and whose longitude is 80° West (of the prime meridian) is shown in Figure 11.74. Assuming that the Earth is spherical and has a radius of 4000 miles, you would label this point as



THE SPHERICAL COORDINATE SYSTEM

In a **spherical coordinate system**, a point *P* in space is represented by an ordered triple (ρ , θ , ϕ).

- **1.** ρ is the distance between *P* and the origin, $\rho \ge 0$.
- **2.** θ is the same angle used in cylindrical coordinates for $r \ge 0$.
- **3.** ϕ is the angle *between* the positive *z*-axis and the line segment \overrightarrow{OP} , $0 \le \phi \le \pi$.

Note that the first and third coordinates, ρ and ϕ , are nonnegative. ρ is the lowercase Greek letter *rho*, and ϕ is the lowercase Greek letter *phi*.



Spherical coordinates Figure 11.75

The relationship between rectangular and spherical coordinates is illustrated in Figure 11.75. To convert from one system to the other, use the following.

Spherical to rectangular:

 $x = \rho \sin \phi \cos \theta, \quad y = \rho \sin \phi \sin \theta, \quad z = \rho \cos \phi$

Rectangular to spherical:

$$\rho^2 = x^2 + y^2 + z^2, \quad \tan \theta = \frac{y}{x}, \quad \phi = \arccos\left(\frac{z}{\sqrt{x^2 + y^2 + z^2}}\right)$$

To change coordinates between the cylindrical and spherical systems, use the following.

Spherical to cylindrical $(r \ge 0)$:

 $r^2 = \rho^2 \sin^2 \phi, \qquad \theta = \theta, \qquad z = \rho \cos \phi$

Cylindrical to spherical $(r \ge 0)$:

$$\rho = \sqrt{r^2 + z^2}, \qquad \theta = \theta, \qquad \phi = \arccos\left(\frac{z}{\sqrt{r^2 + z^2}}\right)$$

The spherical coordinate system is useful primarily for surfaces in space that have a *point* or *center* of symmetry. For example, Figure 11.76 shows three surfaces with simple spherical equations.



EXAMPLE 5 Rectangular-to-Spherical Conversion

Find an equation in spherical coordinates for the surface represented by each rectangular equation.

- **a.** Cone: $x^2 + y^2 = z^2$
- **b.** Sphere: $x^2 + y^2 + z^2 4z = 0$

Solution

a. Making the appropriate replacements for *x*, *y*, and *z* in the given equation yields the following.

$$x^{2} + y^{2} = z^{2}$$

$$\rho^{2} \sin^{2} \phi \cos^{2} \theta + \rho^{2} \sin^{2} \phi \sin^{2} \theta = \rho^{2} \cos^{2} \phi$$

$$\rho^{2} \sin^{2} \phi (\cos^{2} \theta + \sin^{2} \theta) = \rho^{2} \cos^{2} \phi$$

$$\rho^{2} \sin^{2} \phi = \rho^{2} \cos^{2} \phi$$

$$\frac{\sin^{2} \phi}{\cos^{2} \phi} = 1$$

$$\rho \ge 0$$

$$\tan^{2} \phi = 1$$

$$\phi = \pi/4 \text{ or } \phi = 3\pi/4$$

The equation $\phi = \pi/4$ represents the *upper* half-cone, and the equation $\phi = 3\pi/4$ represents the *lower* half-cone.

b. Because $\rho^2 = x^2 + y^2 + z^2$ and $z = \rho \cos \phi$, the given equation has the following spherical form.

 $\rho^2 - 4\rho \cos \phi = 0 \implies \rho(\rho - 4\cos \phi) = 0$

Temporarily discarding the possibility that $\rho = 0$, you have the spherical equation

 $\rho - 4\cos\phi = 0$ or $\rho = 4\cos\phi$.

Note that the solution set for this equation includes a point for which $\rho = 0$, so nothing is lost by discarding the factor ρ . The sphere represented by the equation $\rho = 4 \cos \phi$ is shown in Figure 11.77.





11.7 Exercises

See www.CalcChat.com for worked-out solutions to odd-numbered exercises.

In Exercises 1–6, convert the point from cylindrical coordinates to rectangular coordinates.

1. (-7, 0, 5)2. $(2, -\pi, -4)$ 3. $(3, \pi/4, 1)$ 4. $(6, -\pi/4, 2)$ 5. $(4, 7\pi/6, 3)$ 6. $(-0.5, 4\pi/3, 8)$

In Exercises 7–12, convert the point from rectangular coordinates to cylindrical coordinates.

7. (0, 5, 1)	8. $(2\sqrt{2}, -2\sqrt{2}, 4)$
9. $(2, -2, -4)$	10. $(3, -3, 7)$
11. $(1, \sqrt{3}, 4)$	12. $(2\sqrt{3}, -2, 6)$

In Exercises 13–20, find an equation in cylindrical coordinates for the equation given in rectangular coordinates.

13. $z = 4$	14. $x = 9$
15. $x^2 + y^2 + z^2 = 17$	16. $z = x^2 + y^2 - 11$
17. $y = x^2$	18. $x^2 + y^2 = 8x$
19. $y^2 = 10 - z^2$	20. $x^2 + y^2 + z^2 - 3z = 0$

In Exercises 21–28, find an equation in rectangular coordinates for the equation given in cylindrical coordinates, and sketch its graph.

21.	r = 3	22.	z =	2
23.	$\theta = \pi/6$	24.	r =	$\frac{1}{2}z$
25.	$r^2 + z^2 = 5$	26.	z =	$r^2 \cos^2$
27.	$r = 2 \sin \theta$	28.	r =	2 cos 6

In Exercises 29–34, convert the point from rectangular coordinates to spherical coordinates.

A

29. (4, 0, 0)	30. $(-4, 0, 0)$
31. $(-2, 2\sqrt{3}, 4)$	32. $(2, 2, 4\sqrt{2})$
33. $(\sqrt{3}, 1, 2\sqrt{3})$	34. (-1, 2, 1)

In Exercises 35–40, convert the point from spherical coordinates to rectangular coordinates.

35. $(4, \pi/6, \pi/4)$	36. $(12, 3\pi/4, \pi/9)$
37. $(12, -\pi/4, 0)$	38. $(9, \pi/4, \pi)$
39. $(5, \pi/4, 3\pi/4)$	40. $(6, \pi, \pi/2)$

In Exercises 41–48, find an equation in spherical coordinates for the equation given in rectangular coordinates.

41. $y = 2$	42. <i>z</i> = 6
43. $x^2 + y^2 + z^2 = 49$	44. $x^2 + y^2 - 3z^2 = 0$
45. $x^2 + y^2 = 16$	46. <i>x</i> = 13
47. $x^2 + y^2 = 2z^2$	48. $x^2 + y^2 + z^2 - 9z = 0$

In Exercises 49–56, find an equation in rectangular coordinates for the equation given in spherical coordinates, and sketch its graph.

49. $\rho = 5$	$50. \ \theta = \frac{3\pi}{4}$		
51. $\phi = \frac{\pi}{6}$	52. $\phi = \frac{\pi}{2}$		
53. $\rho = 4 \cos \phi$	54. $\rho = 2 \sec \phi$		
55. $\rho = \csc \phi$	56. $\rho = 4 \csc \phi \sec \theta$		

In Exercises 57–64, convert the point from cylindrical coordinates to spherical coordinates.

57. $(4, \pi/4, 0)$	58. $(3, -\pi/4, 0)$
59. $(4, \pi/2, 4)$	60. $(2, 2\pi/3, -2)$
61. $(4, -\pi/6, 6)$	62. $(-4, \pi/3, 4)$
63. (12, π, 5)	64. $(4, \pi/2, 3)$

In Exercises 65–72, convert the point from spherical coordinates to cylindrical coordinates.

65. $(10, \pi/6, \pi/2)$	66. $(4, \pi/18, \pi/2)$
67. $(36, \pi, \pi/2)$	68. (18, $\pi/3$, $\pi/3$)
69. $(6, -\pi/6, \pi/3)$	70. $(5, -5\pi/6, \pi)$
71. $(8, 7\pi/6, \pi/6)$	72. $(7, \pi/4, 3\pi/4)$

CAS In Exercises 73–88, use a computer algebra system or graphing utility to convert the point from one system to another among the rectangular, cylindrical, and spherical coordinate systems.

	Rectangular	Cylindrical	Spherical
73.	(4, 6, 3)		
74.	(6, -2, -3)		
75.		$(5, \pi/9, 8)$	
76.		(10, -0.75, 6)	
77.			$(20, 2\pi/3, \pi/4)$
78.			(7.5, 0.25, 1)
79.	(3, -2, 2)		
80.	$\left(3\sqrt{2}, 3\sqrt{2}, -3\right)$		
81.	(5/2, 4/3, -3/2)		
82.	(0, -5, 4)		
83.		$(5, 3\pi/4, -5)$	
84.		$(-2, 11\pi/6, 3)$	
85.		(-3.5, 2.5, 6)	
86.		(8.25, 1.3, -4)	
87.			$(3, 3\pi/4, \pi/3)$
88.			$(8, -\pi/6, \pi)$

In Exercises 89–94, match the equation (written in terms of cylindrical or spherical coordinates) with its graph. [The graphs are labeled (a), (b), (c), (d), (e), and (f).]



WRITING ABOUT CONCEPTS

- **95.** Give the equations for the coordinate conversion from rectangular to cylindrical coordinates and vice versa.
- **96.** Explain why in spherical coordinates the graph of $\theta = c$ is a half-plane and not an entire plane.
- **97.** Give the equations for the coordinate conversion from rectangular to spherical coordinates and vice versa.

CAPSTONE

- **98.** (a) For constants *a*, *b*, and *c*, describe the graphs of the equations r = a, $\theta = b$, and z = c in cylindrical coordinates.
 - (b) For constants a, b, and c, describe the graphs of the equations ρ = a, θ = b, and φ = c in spherical coordinates.

In Exercises 99–106, convert the rectangular equation to an equation in (a) cylindrical coordinates and (b) spherical coordinates.

99. $x^2 + y^2 + z^2 = 25$	100. $4(x^2 + y^2) = z^2$
101. $x^2 + y^2 + z^2 - 2z = 0$	102. $x^2 + y^2 = z$
103. $x^2 + y^2 = 4y$	104. $x^2 + y^2 = 36$
105. $x^2 - y^2 = 9$	106. $y = 4$

In Exercises 107–110, sketch the solid that has the given description in cylindrical coordinates.

107. $0 \le \theta \le \pi/2, 0 \le r \le 2, 0 \le z \le 4$ **108.** $-\pi/2 \le \theta \le \pi/2, 0 \le r \le 3, 0 \le z \le r \cos \theta$ **109.** $0 \le \theta \le 2\pi, 0 \le r \le a, r \le z \le a$ **110.** $0 \le \theta \le 2\pi, 2 \le r \le 4, z^2 \le -r^2 + 6r - 8$

In Exercises 111–114, sketch the solid that has the given description in spherical coordinates.

111.
$$0 \le \theta \le 2\pi, 0 \le \phi \le \pi/6, 0 \le \rho \le a \sec \phi$$

112. $0 \le \theta \le 2\pi, \pi/4 \le \phi \le \pi/2, 0 \le \rho \le 1$
113. $0 \le \theta \le \pi/2, 0 \le \phi \le \pi/2, 0 \le \rho \le 2$
114. $0 \le \theta \le \pi, 0 \le \phi \le \pi/2, 1 \le \rho \le 3$

Think About It In Exercises 115–120, find inequalities that describe the solid, and state the coordinate system used. Position the solid on the coordinate system such that the inequalities are as simple as possible.

- 115. A cube with each edge 10 centimeters long
- **116.** A cylindrical shell 8 meters long with an inside diameter of 0.75 meter and an outside diameter of 1.25 meters
- **117.** A spherical shell with inside and outside radii of 4 inches and 6 inches, respectively
- **118.** The solid that remains after a hole 1 inch in diameter is drilled through the center of a sphere 6 inches in diameter
- **119.** The solid inside both $x^2 + y^2 + z^2 = 9$ and

$$\left(x - \frac{3}{2}\right)^2 + y^2 = \frac{9}{4}$$

120. The solid between the spheres $x^2 + y^2 + z^2 = 4$ and $x^2 + y^2 + z^2 = 9$, and inside the cone $z^2 = x^2 + y^2$

True or False? In Exercises 121–124, determine whether the statement is true or false. If it is false, explain why or give an example that shows it is false.

- **121.** In cylindrical coordinates, the equation r = z is a cylinder.
- **122.** The equations $\rho = 2$ and $x^2 + y^2 + z^2 = 4$ represent the same surface.
- **123.** The cylindrical coordinates of a point (x, y, z) are unique.
- **124.** The spherical coordinates of a point (x, y, z) are unique.
- **125.** Identify the curve of intersection of the surfaces (in cylindrical coordinates) $z = \sin \theta$ and r = 1.
- **126.** Identify the curve of intersection of the surfaces (in spherical coordinates) $\rho = 2 \sec \phi$ and $\rho = 4$.

11 REVIEW EXERCISES

See www.CalcChat.com for worked-out solutions to odd-numbered exercises.

In Exercises 1 and 2, let $u = \overline{PQ}$ and $v = \overline{PR}$, and (a) write u and v in component form, (b) write u as the linear combination of the standard unit vectors i and j, (c) find the magnitude of v, and (d) find 2u + v.

1. P = (1, 2), Q = (4, 1), R = (5, 4)**2.** P = (-2, -1), Q = (5, -1), R = (2, 4)

In Exercises 3 and 4, find the component form of v given its magnitude and the angle it makes with the positive *x*-axis.

3.
$$\|\mathbf{v}\| = 8, \ \theta = 60^{\circ}$$
 4. $\|\mathbf{v}\| = \frac{1}{2}, \ \theta = 225^{\circ}$

- **5.** Find the coordinates of the point in the *xy*-plane four units to the right of the *xz*-plane and five units behind the *yz*-plane.
- **6.** Find the coordinates of the point located on the *y*-axis and seven units to the left of the *xz*-plane.

In Exercises 7 and 8, determine the location of a point (x, y, z) that satisfies the condition.

7.
$$yz > 0$$
 8. $xy < 0$

In Exercises 9 and 10, find the standard equation of the sphere.

- **9.** Center: (3, -2, 6); Diameter: 15
- **10.** Endpoints of a diameter: (0, 0, 4), (4, 6, 0)

In Exercises 11 and 12, complete the square to write the equation of the sphere in standard form. Find the center and radius.

11. $x^2 + y^2 + z^2 - 4x - 6y + 4 = 0$ **12.** $x^2 + y^2 + z^2 - 10x + 6y - 4z + 34 = 0$

In Exercises 13 and 14, the initial and terminal points of a vector are given. (a) Sketch the directed line segment, (b) find the component form of the vector, (c) write the vector using standard unit vector notation, and (d) sketch the vector with its initial point at the origin.

13. Initial point: (2, -1, 3)
Terminal point: (4, 4, -7)
14. Initial point: (6, 2, 0)
Terminal point: (3, -3, 8)

In Exercises 15 and 16, use vectors to determine whether the points are collinear.

15. (3, 4, -1), (-1, 6, 9), (5, 3, -6)**16.** (5, -4, 7), (8, -5, 5), (11, 6, 3)

- 17. Find a unit vector in the direction of $\mathbf{u} = \langle 2, 3, 5 \rangle$.
- **18.** Find the vector **v** of magnitude 8 in the direction $\langle 6, -3, 2 \rangle$.

In Exercises 19 and 20, let $u = \overrightarrow{PQ}$ and $v = \overrightarrow{PR}$, and find (a) the component forms of u and v, (b) $u \cdot v$, and (c) $v \cdot v$.

19. P = (5, 0, 0), Q = (4, 4, 0), R = (2, 0, 6)**20.** P = (2, -1, 3), Q = (0, 5, 1), R = (5, 5, 0) In Exercises 21 and 22, determine whether u and v are orthogonal, parallel, or neither.

21.
$$\mathbf{u} = \langle 7, -2, 3 \rangle$$

 $\mathbf{v} = \langle -1, 4, 5 \rangle$
22. $\mathbf{u} = \langle -4, 3, -6 \rangle$
 $\mathbf{v} = \langle 16, -12, 24 \rangle$

In Exercises 23–26, find the angle θ between the vectors.

23.
$$\mathbf{u} = 5[\cos(3\pi/4)\mathbf{i} + \sin(3\pi/4)\mathbf{j}]$$

 $\mathbf{v} = 2[\cos(2\pi/3)\mathbf{i} + \sin(2\pi/3)\mathbf{j}]$
24. $\mathbf{u} = 6\mathbf{i} + 2\mathbf{j} - 3\mathbf{k}, \quad \mathbf{v} = -\mathbf{i} + 5\mathbf{j}$
25. $\mathbf{u} = \langle 10, -5, 15 \rangle, \quad \mathbf{v} = \langle -2, 1, -3 \rangle$
26. $\mathbf{u} = \langle 1, 0, -3 \rangle, \quad \mathbf{v} = \langle 2, -2, 1 \rangle$

- 27. Find two vectors in opposite directions that are orthogonal to the vector $\mathbf{u} = \langle 5, 6, -3 \rangle$.
- **28.** *Work* An object is pulled 8 feet across a floor using a force of 75 pounds. The direction of the force is 30° above the horizontal. Find the work done.

In Exercises 29–38, let $u = \langle 3, -2, 1 \rangle$, $v = \langle 2, -4, -3 \rangle$, and $w = \langle -1, 2, 2 \rangle$.

- **29.** Show that $\mathbf{u} \cdot \mathbf{u} = \|\mathbf{u}\|^2$.
- **30.** Find the angle between **u** and **v**.
- **31.** Determine the projection of **w** onto **u**.
- **32.** Find the work done in moving an object along the vector **u** if the applied force is **w**.
- **33.** Determine a unit vector perpendicular to the plane containing **v** and **w**.
- **34.** Show that $\mathbf{u} \times \mathbf{v} = -(\mathbf{v} \times \mathbf{u})$.
- 35. Find the volume of the solid whose edges are u, v, and w.
- **36.** Show that $\mathbf{u} \times (\mathbf{v} + \mathbf{w}) = (\mathbf{u} \times \mathbf{v}) + (\mathbf{u} \times \mathbf{w})$.
- 37. Find the area of the parallelogram with adjacent sides **u** and **v**.
- 38. Find the area of the triangle with adjacent sides v and w.
- **39.** *Torque* The specifications for a tractor state that the torque on a bolt with head size $\frac{7}{8}$ inch cannot exceed 200 foot-pounds. Determine the maximum force $\|\mathbf{F}\|$ that can be applied to the wrench in the figure.



40. *Volume* Use the triple scalar product to find the volume of the parallelepiped having adjacent edges $\mathbf{u} = 2\mathbf{i} + \mathbf{j}$, $\mathbf{v} = 2\mathbf{j} + \mathbf{k}$, and $\mathbf{w} = -\mathbf{j} + 2\mathbf{k}$.

In Exercises 41 and 42, find sets of (a) parametric equations and (b) symmetric equations of the line through the two points. (For each line, write the direction numbers as integers.)

41. (3, 0, 2), (9, 11, 6) **42.** (-1, 4, 3), (8, 10, 5)

In Exercises 43–46, (a) find a set of parametric equations for the line, (b) find a set of symmetric equations for the line, and (c) sketch a graph of the line.

- **43.** The line passes through the point (1, 2, 3) and is perpendicular to the *xz*-plane.
- **44.** The line passes through the point (1, 2, 3) and is parallel to the line given by x = y = z.
- **45.** The intersection of the planes 3x 3y 7z = -4 and x y + 2z = 3.
- **46.** The line passes through the point (0, 1, 4) and is perpendicular to $\mathbf{u} = \langle 2, -5, 1 \rangle$ and $\mathbf{v} = \langle -3, 1, 4 \rangle$.

In Exercises 47–50, find an equation of the plane and sketch its graph.

47. The plane passes through

(-3, -4, 2), (-3, 4, 1), and (1, 1, -2).

- **48.** The plane passes through the point (-2, 3, 1) and is perpendicular to $\mathbf{n} = 3\mathbf{i} \mathbf{j} + \mathbf{k}$.
- 49. The plane contains the lines given by

 $\frac{x-1}{-2} = y = z + 1$

and

$$\frac{x+1}{-2} = y - 1 = z - 2.$$

- 50. The plane passes through the points (5, 1, 3) and (2, -2, 1) and is perpendicular to the plane 2x + y z = 4.
- **51.** Find the distance between the point (1, 0, 2) and the plane 2x 3y + 6z = 6.
- 52. Find the distance between the point (3, -2, 4) and the plane 2x 5y + z = 10.
- 53. Find the distance between the planes 5x 3y + z = 2 and 5x 3y + z = -3.
- 54. Find the distance between the point (-5, 1, 3) and the line given by x = 1 + t, y = 3 2t, and z = 5 t.

In Exercises 55-64, describe and sketch the surface.

55. x + 2y + 3z = 6 **56.** $y = z^2$ **57.** $y = \frac{1}{2}z$

58. $y = \cos z$

- **59.** $\frac{x^2}{16} + \frac{y^2}{9} + z^2 = 1$ **60.** $16x^2 + 16y^2 - 9z^2 = 0$ **61.** $\frac{x^2}{16} - \frac{y^2}{9} + z^2 = -1$ **62.** $\frac{x^2}{25} + \frac{y^2}{4} - \frac{z^2}{100} = 1$ **63.** $x^2 + z^2 = 4$ **64.** $y^2 + z^2 = 16$
- **65.** Find an equation of a generating curve of the surface of revolution $y^2 + z^2 4x = 0$.
- 66. Find an equation of a generating curve of the surface of revolution $x^2 + 2y^2 + z^2 = 3y$.
- **67.** Find an equation for the surface of revolution generated by revolving the curve $z^2 = 2y$ in the *yz*-plane about the *y*-axis.
- **68.** Find an equation for the surface of revolution generated by revolving the curve 2x + 3z = 1 in the *xz*-plane about the *x*-axis.

In Exercises 69 and 70, convert the point from rectangular coordinates to (a) cylindrical coordinates and (b) spherical coordinates.

69.
$$(-2\sqrt{2}, 2\sqrt{2}, 2)$$
 70. $(\frac{\sqrt{3}}{4}, \frac{3}{4}, \frac{3\sqrt{3}}{2})$

In Exercises 71 and 72, convert the point from cylindrical coordinates to spherical coordinates.

71.
$$\left(100, -\frac{\pi}{6}, 50\right)$$
 72. $\left(81, -\frac{5\pi}{6}, 27\sqrt{3}\right)$

In Exercises 73 and 74, convert the point from spherical coordinates to cylindrical coordinates.

73.
$$\left(25, -\frac{\pi}{4}, \frac{3\pi}{4}\right)$$

74. $\left(12, -\frac{\pi}{2}, \frac{2\pi}{3}\right)$

In Exercises 75 and 76, convert the rectangular equation to an equation in (a) cylindrical coordinates and (b) spherical coordinates.

75.
$$x^2 - y^2 = 2z$$
 76. $x^2 + y^2 + z^2 = 16$

In Exercises 77 and 78, find an equation in rectangular coordinates for the equation given in cylindrical coordinates, and sketch its graph.

77.
$$r = 5 \cos \theta$$
 78. $z = 4$

In Exercises 79 and 80, find an equation in rectangular coordinates for the equation given in spherical coordinates, and sketch its graph.

79.
$$\theta = \frac{\pi}{4}$$
 80. $\rho = 3 \cos \phi$

P.S. PROBLEM SOLVING

1. Using vectors, prove the Law of Sines: If **a**, **b**, and **c** are the three sides of the triangle shown in the figure, then



- **2.** Consider the function $f(x) = \int_0^x \sqrt{t^4 + 1} dt$.
- (a) Use a graphing utility to graph the function on the interval $-2 \le x \le 2$.
 - (b) Find a unit vector parallel to the graph of f at the point (0, 0).
 - (c) Find a unit vector perpendicular to the graph of *f* at the point (0, 0).
 - (d) Find the parametric equations of the tangent line to the graph of f at the point (0, 0).
- **3.** Using vectors, prove that the line segments joining the midpoints of the sides of a parallelogram form a parallelogram (see figure).



4. Using vectors, prove that the diagonals of a rhombus are perpendicular (see figure).



- 5. (a) Find the shortest distance between the point Q(2, 0, 0) and the line determined by the points $P_1(0, 0, 1)$ and $P_2(0, 1, 2)$.
 - (b) Find the shortest distance between the point Q(2, 0, 0) and the line segment joining the points $P_1(0, 0, 1)$ and $P_2(0, 1, 2)$.
- 6. Let P_0 be a point in the plane with normal vector **n**. Describe the set of points *P* in the plane for which $(\mathbf{n} + \overrightarrow{PP_0})$ is orthogonal to $(\mathbf{n} \overrightarrow{PP_0})$.

- 7. (a) Find the volume of the solid bounded below by the paraboloid $z = x^2 + y^2$ and above by the plane z = 1.
 - (b) Find the volume of the solid bounded below by the elliptic

paraboloid
$$z = \frac{x^2}{a^2} + \frac{y^2}{b^2}$$
 and above by the plane $z = k$

where k > 0.

(c) Show that the volume of the solid in part (b) is equal to one-half the product of the area of the base times the altitude, as shown in the figure.



8. (a) Use the disk method to find the volume of the sphere $x^2 + y^2 + z^2 = r^2$.

(b) Find the volume of the ellipsoid $\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1$.

- **9.** Sketch the graph of each equation given in spherical coordinates.
 - (a) $\rho = 2 \sin \phi$
 - (b) $\rho = 2 \cos \phi$
- **10.** Sketch the graph of each equation given in cylindrical coordinates.

(a)
$$r = 2 \cos \theta$$

- (b) $z = r^2 \cos 2\theta$
- 11. Prove the following property of the cross product.

 $(\mathbf{u} \times \mathbf{v}) \times (\mathbf{w} \times \mathbf{z}) = (\mathbf{u} \times \mathbf{v} \cdot \mathbf{z})\mathbf{w} - (\mathbf{u} \times \mathbf{v} \cdot \mathbf{w})\mathbf{z}$

12. Consider the line given by the parametric equations

x = -t + 3, $y = \frac{1}{2}t + 1$, z = 2t - 1

and the point (4, 3, s) for any real number s.

- (a) Write the distance between the point and the line as a function of *s*.
- (b) Use a graphing utility to graph the function in part (a). Use the graph to find the value of *s* such that the distance between the point and the line is minimum.
- (c) Use the *zoom* feature of a graphing utility to zoom out several times on the graph in part (b). Does it appear that the graph has slant asymptotes? Explain. If it appears to have slant asymptotes, find them.

13. A tetherball weighing 1 pound is pulled outward from the pole by a horizontal force **u** until the rope makes an angle of θ degrees with the pole (see figure).

- (a) Determine the resulting tension in the rope and the magnitude of **u** when $\theta = 30^{\circ}$.
- (b) Write the tension *T* in the rope and the magnitude of **u** as functions of θ. Determine the domains of the functions.
- (c) Use a graphing utility to complete the table.

θ	0°	10°	20°	30°	40°	50°	60°
Т							
 u 							

- (d) Use a graphing utility to graph the two functions for $0^{\circ} \le \theta \le 60^{\circ}$.
- (e) Compare T and $\|\mathbf{u}\|$ as θ increases.
- (f) Find (if possible) $\lim_{\theta \to \pi/2^-} T$ and $\lim_{\theta \to \pi/2^-} \|\mathbf{u}\|$. Are the results what you expected? Explain.





Figure for 13



- **14.** A loaded barge is being towed by two tugboats, and the magnitude of the resultant is 6000 pounds directed along the axis of the barge (see figure). Each towline makes an angle of θ degrees with the axis of the barge.
 - (a) Find the tension in the towlines if $\theta = 20^{\circ}$.
 - (b) Write the tension *T* of each line as a function of θ . Determine the domain of the function.
 - (c) Use a graphing utility to complete the table.



- (d) Use a graphing utility to graph the tension function.
- (e) Explain why the tension increases as θ increases.
- **15.** Consider the vectors $\mathbf{u} = \langle \cos \alpha, \sin \alpha, 0 \rangle$ and $\mathbf{v} = \langle \cos \beta, \sin \beta, 0 \rangle$, where $\alpha > \beta$. Find the cross product of the vectors and use the result to prove the identity

 $\sin(\alpha - \beta) = \sin \alpha \cos \beta - \cos \alpha \sin \beta.$

16. Los Angeles is located at 34.05° North latitude and 118.24° West longitude, and Rio de Janeiro, Brazil is located at 22.90° South latitude and 43.23° West longitude (see figure). Assume that Earth is spherical and has a radius of 4000 miles.



- (a) Find the spherical coordinates for the location of each city.
- (b) Find the rectangular coordinates for the location of each city.
- (c) Find the angle (in radians) between the vectors from the center of Earth to the two cities.
- (d) Find the great-circle distance s between the cities. (*Hint:* $s = r\theta$)
- (e) Repeat parts (a)–(d) for the cities of Boston, located at 42.36° North latitude and 71.06° West longitude, and Honolulu, located at 21.31° North latitude and 157.86° West longitude.
- **17.** Consider the plane that passes through the points *P*, *R*, and *S*. Show that the distance from a point *Q* to this plane is

Distance =
$$\frac{|\mathbf{u} \cdot (\mathbf{v} \times \mathbf{w})|}{\|\mathbf{u} \times \mathbf{v}\|}$$

where $\mathbf{u} = \overrightarrow{PR}$, $\mathbf{v} = \overrightarrow{PS}$, and $\mathbf{w} = \overrightarrow{PQ}$.

18. Show that the distance between the parallel planes $ax + by + cz + d_1 = 0$ and $ax + by + cz + d_2 = 0$ is

Distance =
$$\frac{|d_1 - d_2|}{\sqrt{a^2 + b^2 + c^2}}$$
.

- 19. Show that the curve of intersection of the plane z = 2y and the cylinder $x^2 + y^2 = 1$ is an ellipse.
- **20.** Read the article "Tooth Tables: Solution of a Dental Problem by Vector Algebra" by Gary Hosler Meisters in *Mathematics Magazine*. (To view this article, go to the website *www.matharticles.com.*) Then write a paragraph explaining how vectors and vector algebra can be used in the construction of dental inlays.
Answers to Odd-Numbered Exercises







Radius: 5



77. $\overrightarrow{AB} = \langle 1, 2, 3 \rangle$ $\overrightarrow{CD} = \langle 1, 2, 3 \rangle$ $\overrightarrow{BD} = \langle -2, 1, 1 \rangle$ $\overrightarrow{AC} = \langle -2, 1, 1 \rangle$ Because $\overrightarrow{AB} = \overrightarrow{CD}$ and $\overrightarrow{BD} = \overrightarrow{AC}$, the given points form the vertices of a parallelogram. **79.** 0 **81.** $\sqrt{34}$ **83.** $\sqrt{14}$ **85.** (a) $\frac{1}{3}\langle 2, -1, 2 \rangle$ (b) $-\frac{1}{3}\langle 2, -1, 2 \rangle$ **87.** (a) $(1/\sqrt{38})(3, 2, -5)$ (b) $-(1/\sqrt{38})(3, 2, -5)$ **89.** (a)–(d) Answers will vary. (e) $\mathbf{u} + \mathbf{v} = \langle 4, 7.5, -2 \rangle$ $\|\mathbf{u} + \mathbf{v}\| \approx 8.732$ $\|\mathbf{u}\| \approx 5.099$ $\|\mathbf{v}\| \approx 9.014$ **93.** $\langle 0, 10/\sqrt{2}, 10/\sqrt{2} \rangle$ **95.** $\langle 1, -1, \frac{1}{2} \rangle$ **91.** $\pm \frac{7}{2}$ 97. **99.** (2, -1, 2) $(0, \sqrt{3}, 1)$ $\langle 0, \sqrt{3}, \pm 1 \rangle$ **101**. (a) (b) a = 0, a + b = 0, b = 0(c) a = 1, a + b = 2, b = 1(d) Not possible **103.** x_0 is directed distance to yz-plane. y_0 is directed distance to *xz*-plane. z_0 is directed distance to xy-plane. **105.** $(x - x_0)^2 + (y - y_0)^2 + (z - z_0)^2 = r^2$ 107. 0 **109.** (a) $T = 8L/\sqrt{L^2 - 18^2}, L > 18$

(b) L 20 25 30 35 40 45 50 Т 18.4 11.5 10 9.3 9.0 8.7 8.6 30 L = 18 (d) Proof (e) 30 in. (c) T = 8100

111. $(\sqrt{3}/3)(1, 1, 1)$

113. Tension in cable *AB*: 202.919 N Tension in cable *AC*: 157.909 N Tension in cable *AD*: 226.521 N

115.
$$\left(x - \frac{4}{3}\right)^2 + \left(y - 3\right)^2 + \left(z + \frac{1}{3}\right)^2 = \frac{44}{9}$$

Section 11.3 (page 789) **1.** (a) 17 (b) 25 (c) 25 (d) $\langle -17, 85 \rangle$ (e) 34 **3.** (a) -26 (b) 52 (c) 52 (d) $\langle 78, -52 \rangle$ (e) -52**5.** (a) 2 (b) 29 (c) 29 (d) (0, 12, 10) (e) 4 **7.** (a) 1 (b) 6 (c) 6 (d) $\mathbf{i} - \mathbf{k}$ (e) 2 **9.** 20 **11.** $\pi/2$ **13.** $\arccos\left[-1/(5\sqrt{2})\right] \approx 98.1^{\circ}$ **15.** $\arccos(\sqrt{2}/3) \approx 61.9^{\circ}$ **17.** $\arccos(-8\sqrt{13}/65) \approx 116.3^{\circ}$ 23. Neither **19.** Neither **21.** Orthogonal 25. Orthogonal 27. Right triangle; answers will vary. **29.** Acute triangle: answers will vary. **31.** $\cos \alpha = \frac{1}{2}$ **33.** $\cos \alpha = 0$ $\cos\beta = \frac{2}{3}$ $\cos\beta = 3/\sqrt{13}$ $\cos \gamma = -2/\sqrt{13}$ $\cos \gamma = \frac{2}{3}$ **35.** $\alpha \approx 43.3^\circ, \beta \approx 61.0^\circ, \gamma \approx 119.0^\circ$ **37.** $\alpha \approx 100.5^\circ, \beta \approx 24.1^\circ, \gamma \approx 68.6^\circ$ **39.** Magnitude: 124.310 lb $\alpha \approx 29.48^\circ, \beta \approx 61.39^\circ, \gamma \approx 96.53^\circ$ **41.** $\alpha = 90^{\circ}, \beta = 45^{\circ}, \gamma = 45^{\circ}$ **43.** (a) $\langle 2, 8 \rangle$ (b) $\langle 4, -1 \rangle$ **45.** (a) $\left< \frac{5}{2}, \frac{1}{2} \right>$ (b) $\left< -\frac{1}{2}, \frac{5}{2} \right>$ **47.** (a) $\left< -2, 2, 2 \right>$ (b) $\left< 2, 1, 1 \right>$ **49.** (a) $\left< 0, \frac{33}{25}, \frac{44}{25} \right>$ (b) $\left< 2, -\frac{8}{25}, \frac{6}{25} \right>$ 51. See "Definition of Dot Product," page 783. 53. (a) and (b) are defined. (c) and (d) are not defined because it is not possible to find the dot product of a scalar and a vector or to add a scalar to a vector. 55. See Figure 11.29 on page 787. 57. Yes. || 11 · v || ∥v•n ∥

- **59.** \$12,351.25; Total revenue **61.** (a)–(c) Answers will vary.
- 63. Answers will vary. 65. u
- **67.** Answers will vary. Example: $\langle 12, 2 \rangle$ and $\langle -12, -2 \rangle$
- **69.** Answers will vary. Example: (2, 0, 3) and (-2, 0, -3)
- **71.** (a) 8335.1 lb (b) 47,270.8 lb
- **73.** 425 ft-lb **75.** 2900.2 km-N
- 77. False. For example, (1, 1) · (2, 3) = 5 and (1, 1) · (1, 4) = 5, but (2, 3) ≠ (1, 4).
- **79.** $\arccos(1/\sqrt{3}) \approx 54.7^{\circ}$
- **81.** (a) (0, 0), (1, 1)
 - (b) To $y = x^2$ at (1, 1): $\langle \pm \sqrt{5}/5, \pm 2\sqrt{5}/5 \rangle$ To $y = x^{1/3}$ at (1, 1): $\langle \pm 3\sqrt{10}/10, \pm \sqrt{10}/10 \rangle$ To $y = x^2$ at (0, 0): $\langle \pm 1, 0 \rangle$ To $y = x^{1/3}$ at (0, 0): $\langle 0, \pm 1 \rangle$
 - (c) At (1, 1): $\theta = 45^{\circ}$ At (0, 0): $\theta = 90^{\circ}$

83. (a)
$$(-1, 0), (1, 0)$$

(b) To $y = 1 - x^2$ at $(1, 0)$: $\langle \pm \sqrt{5}/5, \pm 2\sqrt{5}/5 \rangle$
To $y = x^2 - 1$ at $(1, 0)$: $\langle \pm \sqrt{5}/5, \pm 2\sqrt{5}/5 \rangle$
To $y = 1 - x^2$ at $(-1, 0)$: $\langle \pm \sqrt{5}/5, \pm 2\sqrt{5}/5 \rangle$
To $y = x^2 - 1$ at $(-1, 0)$: $\langle \pm \sqrt{5}/5, \pm 2\sqrt{5}/5 \rangle$
(c) At $(1, 0)$: $\theta = 53.13^{\circ}$
At $(-1, 0)$: $\theta = 53.13^{\circ}$
85. Proof
87. (a) $\overset{z}{\downarrow}$ (b) $k\sqrt{2}$ (c) 60° (d) 109.5





Section 11.4 (page 798)



5.
$$-\mathbf{j}$$

7. (a)
$$20\mathbf{i} + 10\mathbf{j} - 16\mathbf{k}$$

(b) $-20\mathbf{i} - 10\mathbf{j} + 16\mathbf{k}$
(c) $\mathbf{0}$
11. $\langle 0, 0, 54 \rangle$
13. $\langle -1, -1, -1 \rangle$
15. $\langle -2, 3, -1 \rangle$
17. $\langle \frac{1}{2}, \frac{1}{$

33.
$$\frac{11}{2}$$
 35. $\frac{\sqrt{16,742}}{2}$ **37.** 10 cos 40° \approx 7.66 ft-lb
39. (a) 84 sin θ

- (b) $42\sqrt{2} \approx 59.40$
- (c) $\theta = 90^{\circ}$; This is what should be expected. When $\theta = 90^{\circ}$, the pipe wrench is horizontal.
- **41.** 1 **43.** 6 **45.** 2 **47.** 75
- **49.** At least one of the vectors is the zero vector.
- **51.** See "Definition of Cross Product of Two Vectors in Space," page 792.
- **53.** The magnitude of the cross product will increase by a factor of 4.
- **55.** False. The cross product of two vectors is not defined in a twodimensional coordinate system.
- **57.** False. Let $\mathbf{u} = \langle 1, 0, 0 \rangle$, $\mathbf{v} = \langle 1, 0, 0 \rangle$, and $\mathbf{w} = \langle -1, 0, 0 \rangle$. Then $\mathbf{u} \times \mathbf{v} = \mathbf{u} \times \mathbf{w} = \mathbf{0}$, but $\mathbf{v} \neq \mathbf{w}$.

59-67. Proofs

Section 11.5 (page 807)



- (b) P = (1, 2, 2), Q = (10, -1, 17), PQ = ⟨9, -3, 15⟩
 (There are many correct answers.) The components of the vector and the coefficients of t are proportional because the line is parallel to PQ.
- (c) $\left(-\frac{1}{5}, \frac{12}{5}, 0\right)$, (7, 0, 12), $\left(0, \frac{7}{3}, \frac{1}{3}\right)$
- **3.** (a) Yes (b) No

Parametric
 Symmetric
 Direction

 Equations (a)
 Equations (b)
 Numbers

 5.

$$x = 3t$$
 $\frac{x}{3} = y = \frac{z}{5}$
 $3, 1, 5$
 $y = t$
 $z = 5t$
 $3, 1, 5$

 7.
 $x = -2 + 2t$
 $\frac{x+2}{2} = \frac{y}{4} = \frac{z-3}{-2}$
 $2, 4, -2$
 $y = 4t$
 $z = 3 - 2t$
 $\frac{x-1}{3} = \frac{y}{-2} = \frac{z-1}{1}$
 $3, -2, 1$

 9.
 $x = 1 + 3t$
 $\frac{x-1}{3} = \frac{y}{-2} = \frac{z-1}{1}$
 $3, -2, 1$
 $y = -2t$
 $z = 1 + t$
 $z = 1 + t$
 $z = 1 + t$
 $z = 1 + t$







- **83.** P_1 and P_2 are parallel. **85.** $P_1 = P_4$ and is parallel to P_2 .
- **87.** The planes have intercepts at (*c*, 0, 0), (0, *c*, 0), and (0, 0, *c*) for each value of *c*.
- **89.** If c = 0, z = 0 is the *xy*-plane; If $c \neq 0$, the plane is parallel to the *x*-axis and passes through (0, 0, 0) and (0, 1, -c).
- **91.** (a) $\theta \approx 65.91^{\circ}$ (b) x = 2

$$y = 1 + t$$

$$z = 1 + 2t$$

- **93.** (2, -3, 2); The line does not lie in the plane.
- **95.** Not intersecting **97.** $6\sqrt{14}/7$ **99.** $11\sqrt{6}/6$
- **101.** $2\sqrt{26}/13$ **103.** $27\sqrt{94}/188$ **105.** $\sqrt{2533}/17$
- **107.** $7\sqrt{3}/3$ **109.** $\sqrt{66}/3$

111. Parametric equations: $x = x_1 + at$, $y = y_1 + bt$, and $z = z_1 + ct$ Symmetric equations: $\frac{x - x_1}{a} = \frac{y - y_1}{b} = \frac{z - z_1}{c}$

You need a vector $\mathbf{v} = \langle a, b, c \rangle$ parallel to the line and a point $P(x_1, y_1, z_1)$ on the line.

- **113.** Simultaneously solve the two linear equations representing the planes and substitute the values back into one of the original equations. Then choose a value for *t* and form the corresponding parametric equations for the line of intersection.
- **115.** (a) Parallel if vector $\langle a_1, b_1, c_1 \rangle$ is a scalar multiple of $\langle a_2, b_2, c_2 \rangle$; $\theta = 0$.
 - (b) Perpendicular if $a_1a_2 + b_1b_2 + c_1c_2 = 0$; $\theta = \pi/2$.
- **117.** cbx + acy + abz = abc

119. Sphere:
$$(x - 3)^2 + (y + 2)^2 + (z - 5)^2 = 16$$

121 . (a)	Year	1999	2000	2001	2002
	z (approx.)	6.25	6.05	5.94	5.76
	Year	2003	2004	2005	
	z (approx.)	5.66	5.56	5.56	

The approximations are close to the actual values.

(b) Answers will vary.



73. The Klein bottle does not have both an "inside" and an "outside." It is formed by inserting the small open end through the side of the bottle and making it contiguous with the top of the bottle.



Rectangular

73. (4, 6, 3)

Cylindrical

(7.211, 0.983, 3)

Spherical

(7.810, 0.983, 1.177)



Copyright 2010 Cengage Learning, Inc. All Rights Reserved. May not be copied, scanned, or duplicated, in whole or in part.

- **13.** (a) Tension: $2\sqrt{3}/3 \approx 1.1547$ lb
 - Magnitude of **u**: $\sqrt{3}/3 \approx 0.5774$ lb
 - (b) $T = \sec \theta$; $\|\mathbf{u}\| = \tan \theta$; Domain: $0^{\circ} \le \theta \le 90^{\circ}$ (c)

)	θ	0°	10°		20°		30°	
	Т	1	1.0154		1.0642		1.1547	
	u	0	0.1763		0.3640		0.5774	
	θ	40° 1.3054 0.8391		50° 1.5557 1.1918		60° 2 1.7321		
	Т							
	u							

2.5 (d)

(e) Both are increasing functions.



(f) $\lim_{\theta \to \pi/2^-} T = \infty$ and $\lim_{\theta \to \pi/2^-} \|\mathbf{u}\| = \infty$ Yes. As θ increases, both *T* and $\|\mathbf{u}\|$ increase.

15. $\langle 0, 0, \cos \alpha \sin \beta - \cos \beta \sin \alpha \rangle$; Proof

17.
$$D = \frac{|\overline{PQ} \cdot \mathbf{n}|}{\|\mathbf{n}\|}$$
$$= \frac{|\mathbf{w} \cdot (\mathbf{u} \times \mathbf{v})|}{\|\mathbf{u} \times \mathbf{v}\|} = \frac{|(\mathbf{u} \times \mathbf{v}) \cdot \mathbf{w}|}{\|\mathbf{u} \times \mathbf{v}\|} = \frac{|\mathbf{u} \cdot (\mathbf{v} \times \mathbf{w})|}{\|\mathbf{u} \times \mathbf{v}\|}$$
19. Proof

This page contains answers for this chapter only.